

**PACIFIC SALMON COMMISSION
JOINT CHUM TECHNICAL COMMITTEE
2013 POST SEASON SUMMARY REPORT
TCCHUM (16)-1**

May 2016



MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2013

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Kit Rawson, Co-Chair, Tulalip Tribes
Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Ms. Amy Seiders, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2014

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2015

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO
Ms. Louise de Mestral Bezanson, CDFO

United States Members

Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2016

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

LIST OF ACRONYMS WITH DEFINITIONS

B.C.	British Columbia
CDFO	Canadian Department of Fisheries and Oceans or Fisheries and Oceans Canada
CPUE	Catch per Unit Effort
CU	Conservation Unit (Canada)
CWT	Coded-Wire Tag
ECVI	East Coast Vancouver Island
EO	Economic Opportunity
ESSR	Excess Salmon to Spawning Requirements
ESU	Evolutionarily Significant Unit
FN	First Nation
FN FSC	First Nations Food, Social and Ceremonial
FSC	Food, Social, and Ceremonial (Canada)
GN	Gill net
GSI	Genetic Stock Identification
ITQ	Individual Transferable Quota
mtDNA	Mitochondrial DNA
mSAT	Microsatellite DNA
NOAA	National Oceanic and Atmospheric Administration
NWIFC	Northwest Indian Fisheries Commission
PFMA	Pacific Fishery Management Area
PNPTC	Point No Point Treaty Council
PSC	Pacific Salmon Commission
PST	Pacific Salmon Treaty
SJF	Strait of Juan de Fuca
SJI/PR	San Juan Islands/Point Roberts
SN	Seine net
SNPs	Single-Nucleotide Polymorphisms
TCCHUM	Chum Technical Committee
TR	Troll
U.S.	United States of America
WA	Washington State
WCVI	West Coast Vancouver Island
WDFW	Washington Department of Fish and Wildlife
WSP	Wild Salmon Policy (Canada's Policy for Conservation of Wild Pacific Salmon)

TABLE OF CONTENTS

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2013	i
MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2014	i
MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2015	i
MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2016	i
LIST OF ACRONYMS WITH DEFINITIONS	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	v
LIST OF FIGURES	vi
LIST OF APPENDICES	vi
1 Executive Summary	1
2 Status of Treaty Requirements	1
3 Southern British Columbia Chum salmon	7
3.1 Introduction	7
3.2 Status of Treaty Requirements	7
3.3 Conservation and Harvest Management Strategies	7
3.3.1 Johnstone Strait Chum Salmon Management Strategy	7
3.3.2 Strait of Georgia Chum Salmon Management Strategy	10
3.3.3 Fraser River Chum Salmon Management Strategy	18
3.3.4 West Coast Vancouver Island Chum Salmon Management Strategy	20
3.4 Annual Fishery Descriptions	22
3.4.1 Johnstone Strait	23
3.4.2 Strait of Georgia	24
3.4.3 Fraser River	25
3.4.4 West Coast Vancouver Island	30
3.5 Escapement	31
3.5.1 Inside Southern	31
3.5.2 West Coast Vancouver Island	32

4	United States Chum salmon.....	33
4.1	Washington Run Sizes, Catches, and Spawning Escapements.....	33
4.2	United States Strait of Juan de Fuca Fisheries (Areas 4B, 5, 6C).....	39
4.2.1	Management Intent.....	39
4.2.2	Fishery Description.....	39
4.3	San Juan Islands / Point Roberts Fisheries (Areas 7 and 7A).....	39
4.3.1	Management Intent.....	39
4.3.2	Fishery Description.....	39
5	Stock Identification.....	40
5.1	Tagging of Adult Chum Salmon.....	40
5.2	Otolith, Fin Marking and Coded Wire Tagging (CWT).....	40
5.2.1	Canada.....	40
5.2.2	United States.....	42
5.3	Genetic Stock Identification.....	44
5.3.1	Fishery Sample Collection for Genetic Stock Identification.....	44
5.3.2	Baseline Collection for Genetic Stock Identification.....	45
	REFERENCES CITED.....	47
	APPENDICES.....	48
	Appendix A. ANNEX IV, CHAPTER 6, OF THE PACIFIC SALMON TREATY.....	48
	Appendix B-1. United States commercial salmon catch areas for the Strait of Juan de Fuca and Northern Puget Sound.....	51
	Appendix B-2. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Marine areas.....	52
	Appendix B-3. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Lower Fraser River and surrounding areas.....	53
	Appendix C. PSC Work Plan for the Joint Chum Technical Committee, 2013.....	54
	Appendix D. Concept proposals (see work plans) submitted to Pacific Salmon Commission Southern Endowment Fund.....	56

LIST OF TABLES

Table 3-1. Aggregate Inner South Coast Chum total return, harvest and spawning escapements. Comparison of fixed target and actual harvest rates in the Johnstone Strait fisheries, 2004-2013 ¹	9
Table 3-2. Current spawning escapement goal, hatchery requirements and overall escapement goal for Chum salmon returning to the Strait of Georgia by PFMA's and system.	11
Table 3-3. Key decision points for Fraser River Chum salmon management	19
Table 3-4. Estimated harvest of summer Chum salmon by PFMA for commercial, research and test fishing vessels 1 (July through the second week of September), 2004-2013.	23
Table 3-5. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in Johnstone Strait by PFMA, 2013	24
Table 3-6. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in the Strait of Georgia by PFMA, 2013.	25
Table 3-7. Estimated harvest of Chum salmon from salmon fisheries occurring in Area 29 (both marine and in-river) and in Region 2 (non-tidal) of the Fraser River, 2013.	26
Table 3-8. Estimated harvest of Chum salmon from First Nations and commercial salmon fisheries occurring in Area 29 and Fraser River, 2004-2013	27
Table 3-9. Estimated harvest of Chum salmon in assessed (i.e. surveyed) lower Fraser River recreational fisheries, 2004-2013	29
Table 3-10. Nitinat area harvest, hatchery broodstock collection and spawning escapement estimates by PFMA, 2004-2013.	30
Table 3-11. Inside Southern Chum salmon net escapement estimates (spawning escapement plus hatchery broodstock) for Fraser and Non-Fraser stock aggregates, 2004-2013 ¹ .	31
Table 3-12. WCVI Chum salmon spawning escapement estimates ¹ by PFMA, 2004-2013.	32
Table 4-1. Puget Sound summer Chum salmon pre-and post-season estimates of run size and spawning escapements, 2004-2013	33
Table 4-2. Washington fall Chum salmon pre-and post-season estimates of run size and spawning escapements, 2004-2013	36
Table 4-3. Washington winter Chum salmon pre- and post-season estimates of run size and spawning escapements, 2004-2013	36
Table 4-4. Harvest of Chum salmon in the Strait of Juan de Fuca (SJF) and the San Juan Islands (SJI) commercial fisheries during the summer Chum salmon accounting period, July 1-September 15, 2004-2013.	37
Table 4-5. Harvest of summer, fall, and winter Chum salmon in SJF, SJVPR, Puget Sound and Washington coastal areas, 2004-2013 ¹ .	38

Table 5-1. Releases of Chum salmon with thermally-marked otoliths from WCVI based Canadian hatchery facilities, 2004–2013	41
Table 5-2. Releases of marked Chum salmon from southern B.C. based Canadian hatchery facilities, 2004 to 2013	42
Table 5-3. Numbers of Hood Canal summer Chum salmon released with thermally-marked otoliths or an adipose fin clip by broodyear, 2004–2013 (Point No Point Treaty Tribes and Washington Department of Fish and Wildlife 2014)	43
Table 5-4. Numbers of Puget Sound and Lower Columbia River fall Chum salmon released with thermally-marked otoliths or an adipose fin clip by broodyear, 2004–2013	43
Table 5-5. Chum salmon tissue samples collected in Johnstone Strait fisheries and analyzed for GSI, 2004–2013	44
Table 5-6. Chum salmon tissue samples collected in Areas 7 and 7a analyzed for GSI in 2012–2013	45
Table 5-7. Fall Chum salmon GSI tissue collections from Southern B.C. by Conservation Unit, collection site, life stage, and influence ¹ of enhancement, 2013	45
Table 5-8. Summer, fall and winter Chum salmon GSI tissue collections from W.A. State, 2013	46

LIST OF FIGURES

Figure 2-1. Southern Chum Strategic Plan developed by the joint Chum Technical Committee in 2012	2
Figure 4-1. Puget Sound summer Chum salmon abundance, 2004–2013, by region of origin (JDF: Strait of Juan de Fuca, HCOD: Hood Canal, SPS: South Puget Sound) ... Error! Bookmark not defined.	
Figure 4-2. Puget Sound fall Chum salmon abundance, 2004–2013, by region of origin (JDF: Strait of Juan de Fuca, NOOR: Nookach-Samish, SKAG: Skagit, STSNO: Stillaguamish/Snohomish, SPS: South Puget Sound, HCOD: Hood Canal)	15
Figure 4-3. South Puget Sound winter Chum salmon abundance, 2004–2013	15

LIST OF APPENDICES

Appendix A. ANNEX IV, CHAPTER 6, OF THE PACIFIC SALMON TREATY	48
Appendix B-1. United States commercial salmon catch areas for the Strait of Juan de Fuca and Northern Puget Sound	51
Appendix B-2. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Marine areas	52
Appendix B-3. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Lower Fraser River and surrounding areas	53

1 Executive Summary

This Pacific Salmon Commission (PSC) Joint U.S.-Canada Technical Committee report presents the collaboration on U.S.-Canada science and fisheries in southern British Columbia (BC) and Washington (WA) for the year 2011 to address the specific questions and requirements of Chapter 5, Annex III of the Annexes of the Pacific Salmon Treaty (PST) as follows (Appendix A): The Treaty requires the governments of U.S. and the United States of America (U.S.) concerning Pacific salmon is designed to facilitate co-operation in the management, research and enhancement of Pacific salmon stocks. The U.S.-Canada report, the U.S. and the U.S. requires a Joint U.S.-Canada Technical Committee reporting in the Southern Panel and the U.S. Commission and that certain fisheries the U.S. salmon in southern BC and WA be managed in a specified manner (Appendix A). Certain fisheries of such species, which are specifically mentioned in the PST, are known to have a U.S.-Canada agreement in the other country.

This report presents various aspects of U.S.-Canada science (based in BC) and management (based in WA) and the Southern Panel and the U.S. Commission. This report also discusses the management actions of U.S. and the U.S. in relation to the PST requirements for U.S.-Canada and provides a summary of the key U.S.-Canada science and management information for U.S.-Canada of science in the Treaty. Science in 2011 was slightly better than the recent 5-year average in BC, as slightly above the average in WA. The U.S.-Canada Technical Committee continued with its components of the strategy plan outlined in the 2008 report, which included collecting and exchanging data samples from research vessels and spawning enclosures, conducting genetic sampling workshops in U.S. and Canada, as well as initiating development of a new reconstruction model.

2 Status of Treaty Requirements

U.S.-Canada science and fisheries in southern BC and WA (U.S. Annex III, 1, 2, 3, 4, 5 and 6) (Appendix A) were managed under the terms set out in Chapter 5, Annex III (Appendix A). The following provides a brief overview of the U.S.-Canada and U.S. management actions that were in place in 2011.

Paragraph 1

The Parties agree to maintain a Joint U.S.-Canada Technical Committee (the Committee) reporting under authority agreed in the Southern Panel and the U.S. Commission. Specific tasks for the Committee are listed in Annex III (Chapter 5, Paragraph 1). Some of these include certain stock status, develop and methods for stock management and report on management and research findings.)

The Committee was held two times in 2011. During the PSC Joint Science Meeting in January and during the PSC 12th Annual Meeting in February. In 2011 the Committee developed a strategy plan to guide development and application of a jointly agreed genetic baseline for U.S.-Canada science in southern BC and Washington and a joint development of the U.S.-Canada and the reconstructed World of Salmon (PST) (Figure 1.1). The Committee also completed four proposals for submission to the PSC Southern Fisheries Management and Enhancement Fund (Southern Panel) to carry forward the first three priorities in the strategy plan. Additionally, the

1. **Task 1:** Read the text and answer the questions. (10 points)
The text is about the importance of education in the 21st century. It discusses how technology has changed the way we learn and work. Education is no longer just about memorizing facts, but about developing critical thinking and problem-solving skills. The text also mentions that education is essential for economic growth and social progress.

Question 1:

What is the main topic of the text? (1 point)

- The main topic of the text is the importance of education in the 21st century.
- The main topic of the text is the role of technology in education.
- The main topic of the text is the importance of critical thinking and problem-solving skills.
- The main topic of the text is the importance of education for economic growth and social progress.

2. **Task 2:** Read the text and answer the questions. (10 points)
The text is about the importance of education in the 21st century. It discusses how technology has changed the way we learn and work. Education is no longer just about memorizing facts, but about developing critical thinking and problem-solving skills. The text also mentions that education is essential for economic growth and social progress.

Question 2:

What is the main topic of the text? (1 point)

3. **Task 3:** Read the text and answer the questions. (10 points)
The text is about the importance of education in the 21st century. It discusses how technology has changed the way we learn and work. Education is no longer just about memorizing facts, but about developing critical thinking and problem-solving skills. The text also mentions that education is essential for economic growth and social progress.

Question 3:

What is the main topic of the text? (1 point)

4. **Task 4:** Read the text and answer the questions. (10 points)
The text is about the importance of education in the 21st century. It discusses how technology has changed the way we learn and work. Education is no longer just about memorizing facts, but about developing critical thinking and problem-solving skills. The text also mentions that education is essential for economic growth and social progress.

Question 4:

What is the main topic of the text? (1 point)

- The main topic of the text is the importance of education in the 21st century.
- The main topic of the text is the role of technology in education.
- The main topic of the text is the importance of critical thinking and problem-solving skills.
- The main topic of the text is the importance of education for economic growth and social progress.

catches for the purpose of genetic stock identification sampling shall not be included in the aforementioned limit;

- c) For run sizes above the critical threshold, the base catch ceiling for the U.S. Chum salmon fisheries in Areas 7 and 7A will be 130,000 Chum salmon;*
- d) Canada will provide a run size estimate of Chum salmon entering the Fraser River no later than October 22. If the estimate is less than 300,000, the U.S. will limit its fishery impacts on Fraser River Chum salmon by restricting catch in Areas 7 and 7A to not exceed 20,000 additional Chum salmon from the day following the date the U.S. is notified. The total catch is not to exceed the catch ceiling of 130,000 Chum salmon;*
- e) U.S. commercial fisheries for fall Chum salmon in Areas 7 and 7A will not occur prior to October 30;*
- f) The U.S. will manage the Areas 7 and 7A fisheries for Chum salmon with the intent to minimize the harvest of non-targeted species;*
- g) No U.S. catch shortfalls may be accrued; however any overages shall be carried forward as indicated in (h) and (i);*
- h) Due to management improvisation, a catch in the U.S. of up to 133,000 Chum salmon will not result in an overage calculation. Catches in excess of 133,000 Chum salmon shall result in an overage being calculated by subtracting 130,000 from the total Chum catch. Overages will be accounted for by reducing the U.S. annual catch ceiling in up to two subsequent non-critical Inside Southern Chum salmon years; and*
- i) From the day following the date the U.S. is notified of a run size below the critical threshold as defined in (d)(i) or (d), any catches in excess of 20,000 Chum salmon will result in an overage. Overages will be accounted for by reducing the U.S. annual catch ceilings in up to two subsequent non-critical Inside Southern Chum salmon years.*

In 2003, U.S. fisheries in Areas 7 and 7A were managed in accordance with the above provisions. U.S. commercial fisheries were initiated as scheduled on October 10. An estimated Fraser River Chum salmon run size was provided by Canada prior to October 22, consistent with the Treaty obligations. The fishery was therefore continued without restriction through November 5.

Paragraph 11

The United States shall conduct its Chum salmon fishery in the Strait of Juan de Fuca (United States Areas 4B, 5, and 6C) so as to maintain the limited effort nature of this fishery, and, to the extent practicable, not increase interceptions of Canadian origin Chum salmon. The United States shall continue to monitor this fishery to determine if recent catch levels indicate an increasing level of interception.

This fishery is restricted to Treaty Indian fishers from four tribes and to gill net gear only. Due to the limited effort of the Juan de Fuca fishery, the Committee has not sampled the fishery for contribution of Canadian origin Chum salmon since 1996. Genetic stock identification (GSI)

samples collected from this fishery in prior years indicate the majority of the catch is Chum salmon of U.S. origin (Beattie et al. 1996).

Paragraph 12.

All information concerning by-catch of other salmon species from the Chum salmon fisheries covered by this chapter will be shared between the Parties in the annual Post Season Report.

By-catch information for other salmon species was shared in accordance with the Treaty through each Party's post-season reporting process.

Paragraph 13

Should circumstances arise that are inconsistent with either Parties understanding of the intent of the chapter, the Southern Panel will discuss the matter postseason and explore options for taking the appropriate corrective actions.

No such circumstances arose in 2013.

3 Southern British Columbia Chum salmon

3.1 Introduction

Southern B.C. Chum salmon stocks and fishing areas (Pacific Fishery Management Areas or PFMAAs; Appendix B) are, for the purposes of management, analysis and reporting, grouped into four major units: Johnstone Strait, Strait of Georgia, Fraser River, and the West Coast Vancouver Island (WCVI). Additionally, Chum salmon originating from Johnstone Strait, the Strait of Georgia, and the Fraser River are collectively described as Inside Southern Chum salmon.

3.2 Status of Treaty Requirements

During 2013, the Southern B.C. Chum salmon fisheries were managed according to the requirements of Annex IV, Chapter 6, as amended in 2009 (Appendix A).

3.3 Conservation and Harvest Management Strategies

The general management approach adopted by Fisheries and Oceans Canada (DFO) for Southern B.C. Chum salmon is to achieve wild escapement targets, while augmenting production through enhancement of selected stocks. In practice, this approach is achieved through the application, in mixed stock fisheries, of harvest rates which are compatible with wild or natural stock productivity. If there are Chum salmon stocks that return to their area of origin in numbers above that area's escapement goal, they may be subject to additional harvesting within their terminal areas.

The following describes the management strategies and fishing plans for the four units of Southern B.C. Chum salmon stocks.

3.3.1 Johnstone Strait Chum Salmon Management Strategy

Since 2002, the mixed stock Chum directed fisheries in Johnstone Strait have been managed to approximate a fixed exploitation rate (~20%). Tagging studies conducted from 2000 through 2002 helped in the development of this fixed exploitation rate strategy by assessing the migration timing and harvest rate by fishing gear on an available abundance of Chum salmon in Johnstone Strait. This fixed exploitation strategy continued to be employed through 2013 (DFO 2013).

Some of the key objectives of this strategy are to ensure sufficient escapement levels while providing more stable fishing opportunities (DFO 2013). The exploitation rate is set at 20% across all harvesters, when abundance is estimated to be above a critical level. The critical level, or threshold, was determined to be 1.0 million Chum salmon (Appendix A). The Johnstone Strait Chum salmon test fishery, as well as the information gained from any commercial openings, is used in-season to determine the likelihood of achieving the critical level for Inside Southern Chum salmon. When expected levels are less than 1.0 million, only assessment fisheries and non-commercial fisheries will be conducted and any planned commercial fisheries targeting Chum salmon will be suspended (Appendix A). When the critical level is expected to be achieved, fisheries with an exploitation rate of up to 20% in Johnstone Strait can be conducted (Appendix A).

Of this 20%, the commercial sector is allocated 15%. The remaining 5% is set aside to provide for First Nations' food, social and ceremonial needs (FSC) satisfy recreational and test fishing

requirements, and to provide a buffer to the commercial exploitation (CDFO 2013). The impact of the Johnstone Strait fisheries during fixed exploitation approach (2004–2013), on Inside Southern Chum salmon stocks are detailed in Table 3-1.

The specific objectives of the fixed exploitation rate strategy are to:

- continue to rebuild/maintain stocks to the optimum wild escapement objective (defined as 2.5 million wild Inside Southern Chum salmon);
- establish a pre-season fishing plan to achieve an exploitation rate of 20% on Inside Southern Chum salmon in the Johnstone Strait Chum salmon fishery; and,
- stabilize commercial catch rates to provide opportunities at both low and high abundance levels

Table 3-1. Aggregate Inner South Coast Chum total return, harvest and spawning escapements. Comparison of fixed target and actual harvest rates in the Johnstone Strait fisheries. 2004–2013¹

	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
PRE-SEASON										
Inside Southern Abundance Forecast (or Outlook)	Average	Average	Average	Average to Above Average	Below Average	Below Average	Average to Below Average	Average to Below Average	Average to Below Average	Average to Below Average
POST-SEASON										
Inside Southern Abundance²	5,519,205	3,126,617	4,301,780	2,728,020	2,227,630	2,227,110	1,181,528	3,449,783	3,518,607	2,931,714
Inside Southern Harvest	1,876,239	1,304,237	1,527,453	796,239	579,756	747,043	127,967	1,441,680	911,427	981,945
Est. Inside Southern Harvest Rate	34.0%	41.7%	35.5%	29.2%	26.0%	33.5%	10.8%	41.8%	25.9%	33.49%
Johnstone Strait Harvest										
Commercial, Sport Area 11-13³	1,176,611	856,029	859,201	480,620	299,775	513,884	55,520	750,436	392,608	635,814
First Nations Area 11-13	27,041	18,841	16,453	14,870	11,630	13,576	4,060	9,880	13,106	13,444
Johnstone Strait Harvest Total	1,203,652	874,870	875,654	495,490	311,405	527,460	59,580	760,316	405,714	649,258
Target Johnstone Strait Harvest Rate	20%	20%	20%	20%	20%	20%	20%	20%	20%	20%
Est. Johnstone Strait Harvest Rate	21.8%	28.0%	20.4%	18.2%	14.0%	23.7%	5.0%	22.0%	11.5%	22.2%
ESCAPEMENT (Includes wild and enhanced)⁴										
Inside Southern Escapement	3,642,966	1,822,380	2,774,327	1,931,781	1,647,874	1,480,067	1,053,561	2,008,103	2,607,180	1,949,769

(1) Historic data in this table have been updated with most recent estimates; values may deviate from past reports.

(2) Total Inside Southern abundance includes total Inside Southern harvest plus escapement. Harvest composition based on historic GSI for all fisheries.

(3) Includes commercial, sport and test fishery harvest.

(4) Escapement estimates do not include any removals associated with hatchery rack (e.g. ESSR); those values are included in harvest. Escapement estimates have not been expanded for populations not monitored for escapement

3.3.2 Strait of Georgia Chum Salmon Management Strategy

Chum salmon stocks return to all areas of the Strait of Georgia (mid-Vancouver Island PFMA's 14-19) and are of enhanced and wild (natural) origin. Fisheries can target enhanced returns (e.g. Area 14: Puntledge, Big Qualicum and Little Qualicum rivers), wild returns (e.g. Area 16: main river systems of Tzoonie, Deserter and Skwawka rivers in the Jervis inlet area) and mixed returns (e.g. Area 19: directed primarily at enhanced Goldstream River population although some natural Cowichan River Chum salmon are also harvested).

With one exception, pre-season forecasts of the return of Strait of Georgia Chum salmon for population aggregates corresponding to PFMA's have limited use due to their inability to reasonably predict returns (CDFO 2013). The one exception is Chum salmon returning to Area 14. As a result, unlike some other units of Southern BC Chum Salmon (e.g. WCVI see section 3.3.4), pre-season fishing plans based on a pre-season forecast are not typically developed for Strait of Georgia Chum salmon stocks.

Strait of Georgia Chum salmon are managed as a component of a "mixed-stock harvest strategy" for Johnstone Strait and the northern Strait of Georgia. In-season management of Strait of Georgia Chum salmon is guided by advice from the South Coast Chum Advisory Committee (Chum Advisory Committee or Chum Working Group). The Chum Working Group represents interests for Strait of Georgia (mid-Vancouver Island Areas 14-19), Johnstone Strait, WCVI and Fraser River Chum salmon-directed fisheries. Fishing opportunities for Chum salmon are evaluated at weekly meetings of the Chum Working Group that usually start in the first week of October. In-season data is reviewed on a weekly basis until the conclusion of the fishing season that usually occurs around the end of November. The Chum Working Group includes internal (CDFO staff) and external (First Nations, commercial and recreational) representation and has been in operation since 2004. Additionally, Chum salmon management in Area 18 is also guided by advice from the Cowichan Fisheries Roundtable.

Chum salmon fishing opportunities in the Strait of Georgia PFMA's are predominately shaped by in-season escapement information, test fishing data, commercial harvesting data from Johnstone Strait fisheries when available, and coast-wide allocations of Chum salmon. When escapement goals are achieved (Table 3-2) or when stock abundance allows, commercial gill net, purse seine and troll fisheries can be permitted. As mentioned, the exception is Area 14 where a pre-season forecast of returning Chum abundance is used in addition to the in-season information listed above, to guide management decisions on fishing opportunities (CDFO 2013).

Chum salmon stocks return to all areas of the Strait of Georgia (mid-Vancouver Island PFMA's 14-19) and are of enhanced and wild (natural) origin. Fisheries can target enhanced returns (e.g. Area 14: Puntledge, Big Qualicum and Little Qualicum rivers), wild returns (e.g. Area 16: main river systems of Tzoonie, Deserter and Skwawka rivers in the Jervis inlet area) and mixed returns (e.g. Area 19: directed primarily at enhanced Goldstream River population, although some natural Cowichan River Chum salmon are also harvested).

Table 3-2. Current spawning escapement goal, hatchery requirements and overall escapement goal for Chum salmon returning to the Strait of Georgia by PFMA's and system.

PFMA	System	Spawning Escapement Goal	Hatchery Requirements	Overall Escapement Goal
Area 14	Puntledge River	60,000	10,000	700,000
	Big Qualicum River	100,000		
	Little Qualicum River	130,000		
Area 16	Jervis Inlet Streams	n/a	n/a	120,000
Area 17	Nanaimo River	n/a	n/a	60,000
Area 18	Cowichan River	n/a	n/a	100,000
Area 19	Goldstream River	n/a	n/a	15,000

(n/a) not applicable.

Fishing opportunities in the Strait of Georgia are constrained by concerns over bycatch of passing Chum salmon stocks and local Chinook and Coho salmon populations. These concerns are addressed by Area specific pre- and in-season management measures that may include:

- *area closures*: beach boundaries and limiting fisheries to terminal areas to minimize impacts on passing stocks; and,
- *non-retention*: of Chinook and Coho salmon.

As well, to reduce bycatch mortality and to improve the quality of catch data, additional pre-and in-season management measures may be implemented in the Strait of Georgia fisheries. These additional measures may include:

- *daylight fishing only*: in the gill net fishery;
- *maximum soak time*: in the gill net fishery;
- *mandatory bruiling*: in the seine fishery;
- *barbless hooks*: in the troll fishery; and,
- *on-board observers*: if high bycatch occurs.

Pre- and in-season management measures aimed at reducing bycatch mortality are commonly implemented after consultation with the Chum Working Group.

First Nations FSC fisheries remain a priority and occur in terminal areas targeting individual populations from river systems within or adjacent to band territories. Harvest limits may be set when there are conservation concerns for the target population; however, First Nations FSC fisheries have priority over other fisheries. Generally, catch and effort in First Nations FSC Chum salmon fisheries is low compared to commercial Chum salmon fisheries.

Although tidal recreational fisheries are open throughout the Strait of Georgia, Chum salmon are not commonly targeted by recreational anglers. Nevertheless, the Strait of Georgia recreational Chum salmon catch is estimated annually through a Creel Survey conducted by CDFO. In most years, the Creel Survey operates from the beginning of May through to the end of September. Data from this survey is compiled and analyzed to produce catch and effort statistics by Area for

each salmon species. Non-tidal recreational fisheries vary by Area and are considered if escapement and FSC needs have been met.

3.3.2.1 Area 14 Chum Salmon Management Strategy

Chum salmon returning to Area 14 have been enhanced since the late 1980's, and terminal fisheries targeting the enhanced Chum salmon populations of Portledge, Big Qualicum and Little Qualicum rivers have occurred in October and November since the early 1990's. Returning Area 14 Chum salmon abundance is forecasted pre-season using brood escapement, average survival, and age composition. In-season run strength is assessed from any early catch information, visual observations at river estuaries, and escapement counts to the three enhanced river systems.

The spawning escapement goals for Chum salmon to the three river systems are 80,000 to Portledge River, 100,000 to Big Qualicum River, and 130,000 to Little Qualicum River, including enhancement facility requirements of 10,000, the overall escapement goal is 300,000 Chum salmon for Area 14 (Table 3-2). Spawning escapements are monitored by CDFO Stock Assessment and hatchery staff.

The recommended approach regarding Chum salmon-directed fishing opportunities in Area 14 is made at the first meeting of the Chum Working Group. This meeting is tentatively scheduled for the first week of October. Chum Working Group meetings are subsequently held every week until late November.

Commercial Chum salmon-directed fisheries have a specific harvest strategy, implemented since 1981 that consists of limited early harvest prior to any escapement occurring. The allowable early Chum salmon harvest is set at 65% of the predicted surplus. The predicted surplus is calculated by subtracting the overall escapement goal (300,000, Table 3-2) and a buffer (100,000) from the pre-season terminal run size forecast (CDFO 2011). The buffer safeguards against errors in the pre-season forecast of Area 14 stock abundance.

The limited early commercial harvest of Chum salmon is planned pre-season and executed in-season in the second or third week of October based on whether a surplus, using the formula described above, is identified. However, if catches in the commercial Chum salmon fisheries in Johnstone Strait indicate low Chum salmon abundance, a decision on the approach in Area 14 may be deferred until the following week.

The management objectives for Area 14 are:

- achieve Area 14 Chum salmon escapement requirements of 300,000;
- ensure adequate Chinook and Coho salmon escapements to Area 14 enhancement facilities;
- provide access to First Nations for FSC purposes;
- minimize the harvest of passing salmon populations;
- maximize economic return;
- work towards south coast Chum salmon allocation targets for gill net, seine net and troll sectors;

- attempt to manage wild fisheries in Area 14 as good large salmon (e.g. greater than 100/1000).

Starting in the second or third week of October, opportunities for gill net, seine net and trawl fisheries are based on pre-season forecasts, in-season catch per unit effort (CPUE) information from commercial Chinook salmon fisheries in Johnsons Strait, and management information. Equipment information becomes increasingly important when considering further commercial opportunities in the latter part of October. These commercial opportunities are discussed at the weekly Chinook Working Group meetings and may include consideration for:

- *limited effort seine net fishery*: a limited effort seine fishery with a harvest target will be considered from late October to late November, based on Chinook salmon management, disturbance in the approach area and beach Chinook education guidelines. Full fleet opportunities may also be available;
- *expanded gill net and trawl fishery*: if gear counts indicate a median fleet size of 70 vessels or less, gill net and trawl openings may be expanded beyond one to two days per week, as well, additional fishing opportunities for gill net and trawl may be considered following the seine net fishery, and;
- *additional fishing days*: additional fishing days are considered if fishing time is lost due to poor weather conditions.

Once spawning management goals and hatchery requirements have been achieved, any returns above the overall management goal may be harvested at 100% CPUE.

Commercial fishing opportunities are constrained by concerns over local and passing Chinook, Coho and Chinook salmon populations. Management measures to address these concerns can include:

- *area closures*: beach boundaries to protect Chinook and Coho salmon distributions may range from half a mile to one and a half miles depending upon beach concerns and time of year; French Creek Indian boundary and Baynes Sound area closures to protect wild Chinook and Coho salmon populations;
- *non-retention* of Coho salmon;
- *maximum soak time* in the gill net fishery;
- *mandatory hauling* in the seine fishery;
- *hatchery bands* in the trawl fishery;
- *daylight fishing only* in the gill net fishery if there are significant levels of bycatch; and
- *seasonal boundaries*: designed to minimize impacts on passing salmon populations.

Five Nations FMC fisheries are conducted in Area 14 and at the enhancement facilities prior to consideration of ENR fisheries targeting Chinook salmon. Valid recreational fisheries are subject to the normal daily and possession limits for Chinook salmon (daily limit four per the possession limit eight) and are open throughout the Area.

These arrangements have been confirmed, notwithstanding reduction in C-Team salaries in the Rushcliffe and Big Chalfont rivers, will be considered. These fishing opportunities may consist of up to the second or third week of October and on board, in part, open to women and joint access fishing of C-Team salmon.

C-Team salmon returning in Year 16 have been enhanced since the year 1997, with several fisheries targeting the enhanced C-Team salmon populations of Rushcliffe, Big Chalfont and Little Chalfont rivers have occurred in October and November since the year 1997. However, Year 16 C-Team salmon abundance is forecast to be open entry based on previous average survival and egg-comparisons. To ensure not through a seasonal frost-free, open entry information, visual observations of river education, and management needs in the River enhanced river systems.

3.3.3.3 River 16 C-Team Salmon Management Strategy

This feature targets wild C-Team salmon stocks returning in river systems in the River 16 area. The main systems are Truro, Stour and River 16. The overall management goal for River 16 is to ensure a 100,000. This overall feature will also be included in continued management goals have been agreed.

Year 16 C-Team salmon are managed as a component of "wild stock harvest strategy" for C-Team salmon, and fishing opportunities are provided to work with objectives of C-Team salmon. C-Team salmon fishing opportunities are provided in weekly meetings of the C-Team salmon. C-Team salmon, weekly meeting in the first week of October. Management in the year is controlled by Fisheries and Marine C-Team (Charles River) weekly, 1997-1998, management and fishing fishing River 16. C-Team salmon is provided as a weekly basis with the end of the season, which weekly occurs around the end of November. Year 16 C-Team salmon are not allowed to be open season through down. Fishing opportunities will be provided to open after the management goal has been achieved. Achievement of the management goal includes the number of fish in river plus the amount of fish made a designated minimum goal. The overall potential fishing opportunity is anticipated over the end of October of 1997-1998.

C-Team salmon fishing opportunities are considered by various river fishing groups in C-Team salmon and general C-Team salmon populations. To address these concerns, there is a management and research of C-Team salmon and fishing is based on several goals.

Specifically, C-Team salmon fishing opportunities in Year 16 do not meet as a regular basis. There have been no fishing in Year 16 in recent years.

Year 16 C-Team salmon directed fishing target wild C-Team salmon populations returning in river systems in the River 16 area. The main systems are Truro, Stour and River 16.

3.3.3.4 River 17 C-Team Salmon Management Strategy

C-Team salmon fishing in Year 17 are directed primarily at the Truro River population. The open entry zone, the Truro River C-Team salmon stock is supplemented by production from the Truro River fisheries.

The overall management goal for the Truro River is 40,000 C-Team salmon (Table 3.3). C-Team salmon management estimates are derived from joint 1997/1998 management in river management.

The second dimension of the ethical decision-making process is the ethical approach. The literature is split concerning whether an individual's ethical approach is stable (e.g., Treviño and Youngblood, 1990) or malleable (e.g., Treviño and Weaver, 1999). The latter view is supported by the research that has not only shown that ethical approaches can be learned (e.g., Treviño and Weaver, 1999) but also that they can be changed through the socialization process (e.g., Treviño and Weaver, 1999). The latter view is supported by the research that has shown that ethical approaches can be learned (e.g., Treviño and Weaver, 1999) but also that they can be changed through the socialization process (e.g., Treviño and Weaver, 1999).

Approaches to ethical decision-making are often categorized into two groups: rule-based and principle-based. Rule-based approaches are based on the idea that ethical decisions should be made by applying a set of rules or principles. Principle-based approaches are based on the idea that ethical decisions should be made by applying a set of principles or values.

1. Rule-based approaches are based on the idea that ethical decisions should be made by applying a set of rules or principles.
2. Principle-based approaches are based on the idea that ethical decisions should be made by applying a set of principles or values.
3. Rule-based approaches are based on the idea that ethical decisions should be made by applying a set of rules or principles.
4. Principle-based approaches are based on the idea that ethical decisions should be made by applying a set of principles or values.
5. Rule-based approaches are based on the idea that ethical decisions should be made by applying a set of rules or principles.
6. Principle-based approaches are based on the idea that ethical decisions should be made by applying a set of principles or values.

The research on ethical decision-making is still in its early stages. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.

1. The research on ethical decision-making is still in its early stages.
2. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.
3. The research on ethical decision-making is still in its early stages.
4. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.
5. The research on ethical decision-making is still in its early stages.
6. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.

The research on ethical decision-making is still in its early stages. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.

The research on ethical decision-making is still in its early stages. However, the research that has been conducted to date suggests that ethical decision-making is a complex process that is influenced by a number of factors, including the individual's ethical approach, the situation, and the social context.

- [illegible]

10. **Answer: A** – The passage states that the author is a member of the same organization as the person who wrote the letter. The author is a member of the same organization as the person who wrote the letter. The author is a member of the same organization as the person who wrote the letter.

- *harpoon hooks* in the troll fishery;
- *daylight fishing only* in the gill net fishery if Coho salmon encounters are high;
- *selective fishing techniques* commercial fisheries will utilize selective fishing techniques to minimize bycatch impacts; and,
- *on-board observers* on-board observers could be employed if high bycatch occurs.

2.3.3 Fraser River Chum Salmon Management Strategy

The management strategy for Fraser River Chum salmon includes management goals and harvest allocation objectives for Chum salmon fisheries occurring within the Fraser River. Fisheries are structured to ensure a spawning escapement of 800,000 Chum salmon using in-season information derived from the CDFD Albion Chum salmon test fishery (see section 2.3.3.1). Decision rules that guide the management of Fraser River Chum salmon fisheries are summarized in Table 2-5.

Additionally, management of in-river Chum salmon fisheries is guided by the need to minimize impacts on co-migrating stocks of concern, including Interior Fraser River Coho salmon and Interior Fraser River Steelhead. In order to limit incidental impacts on Interior Fraser River Coho salmon, fishing for salmon with non-selective gear is restricted for all sectors (First Nations, recreational, and commercial) from early September to mid-October within the Fraser River mainstem below Spuzzan Creek. Conservation measures taken to protect Interior Fraser River Steelhead are developed in conjunction with the B.C. Ministry of Environment on an annual basis, and are primarily focused on reducing the impact of Fraser River commercial gill net fisheries. In recent years, these restrictions have resulted in the delay of commercial fishing opportunities for Chum salmon until late October, with the exception of those fisheries that operate with selective gear.

Table 3-3. Key decision points for Fraser River Chum salmon management.

Run Size	Harvest Plan	First Nations	Commercial	Recreational
< 500,000 in Fraser	< 10%	Limited (reduced hours and days/week fishing)	Closed	Main stem Fraser River closed, restricted openings on tributaries
500,000 to 800,000 in Fraser	Directed fisheries limited to FSC	Normal (72,000)	Closed	Main stem Fraser River closed, restricted openings on tributaries
800,000 to 950,000 in Fraser	Catch out to exceed \$1,000 (72,000 First Nations and 9,000 test fishing)	Normal (72,000)	Closed	Main stem Fraser River open, restricted openings on tributaries
950,000 to 1,050,000 in Fraser	Commercial catch out to exceed 10% for Chum	Normal (72,000)	Open (15,000-105,000)	Open
> 1,050,000 in Fraser	Commercial catch out to exceed 15% for Chum	Normal (72,000)	Open	Open

3.3.3.1 Fraser River In-season Terminal Abundance Estimation

Terminal abundance of Fraser River Chum salmon is estimated using a Bayesian model that incorporates pre-season information on run size and migration timing, with in-season information on Chum salmon catch from the Albion Chum salmon test fishery (Garey and Palermo 2000).

The Albion Chum salmon test fishery has operated annually since 1979 on the lower Fraser River in Area 29 near Fort Langley. The test fishery is conducted with a drifted gill net at a specific site near the old Albion ferry crossing. The test fishery begins in early September of each year, and usually fishes until the end of November. On each day of operation, the boat fishes two sets, timed to coincide with the daily high tide. The Albion Chum salmon test fishery normally fishes every other day from September 1st through October 20th, alternating days with the Albion Chinook salmon test fishery (which fishes an 8" mesh gill net during this period). From October 21st through the end of November, the Chum salmon test fishery operates daily. The gill net used in the Albion Chum salmon test fishery is 150 fathoms long, constructed from uniform 6-75" mesh.

The first in-season estimate of terminal Fraser River Chum salmon abundance is typically provided in mid-October. Limited First Nations FSC fisheries for Fraser Chum salmon may be permitted prior to this date, unless a conservation concern has been identified.

3.3.4 West Coast Vancouver Island Chum Salmon Management Strategy

Chum salmon stocks return to all areas on the West Coast of Vancouver Island (WCVI). When stock abundance allows, commercial Chum salmon fisheries are conducted in the marine waters outside Nitinat Lake (Area 21), Barkley Sound (Area 23), Clayoquot Sound (Area 24) and Outer Nootka Sound/Tlupana and Esperanza Inlet (Areas 25). Both commercial gill net and purse seine fleets target WCVI Chum salmon in these Areas.

A pre-season forecast of the return of WCVI Chum salmon is predicted for population aggregates corresponding to CDFO's Pacific Fishery Management Areas. Natural and enhanced returns are approximated through separate models and summed to produce a total return forecast for each Area. However, the relative inaccuracy of the pre-season forecasts resulted in the adoption of a fixed harvest rate strategy in the early 2000's for those fisheries targeting mixed stock aggregates.

With the exception of near-terminal areas, such as Nitinat (Area 21 & 22) or Tlupana Inlet (Area 25), where hatchery stocks dominate, WCVI Chum salmon fisheries are currently managed to an exploitation rate of approximately 20% or less. Exploitation is controlled by limiting fishing effort to specific areas, specific times or to a specific number of vessel fishing days per week. Since 2012, revised fishery lower and target reference points (FLRPs and FTRPs) were developed based on the Sustainable Escapement Goal (SEG) method. The SEG approach uses the long-term escapement series to set fishery reference points. Conservative "SEGs" are defined as the 25% and 75% of a long-term escapement time series. Although the WCVI Chum forecast is highly uncertain, the forecast is used to inform pre-season fishery planning. Where the forecast is below the FLRP for an area, fisheries are curtailed. Where the forecast is below the FTRP, fisheries are more limited (CDFO 2013)

Purse seine opportunities generally occur when excess salmon to spawning requirements (ESSR) are identified in near terminal areas (e.g. hatchery stocks returning to Nitinat River or Tlupana Inlet).

WCVI First Nations FSC fisheries remain a priority and occur in terminal areas targeting individual populations from river systems within or adjacent to band territories. Harvest limits may be set when there are conservation concerns for the target population; however, First Nations FSC fisheries have priority over other fisheries. Generally, harvest and effort in First Nations FSC Chum salmon fisheries is low compared to commercial Chum salmon fisheries.

Chum salmon are not commonly targeted by recreational anglers on the WCVI. Nevertheless, the WCVI recreational Chum salmon catch is estimated annually through a Creel Survey conducted by CDFO. Creel Survey information is augmented by logbook and manifest records of catch and effort submitted by lodges operating guided fishing trips on the WCVI. These data are compiled and analyzed to produce catch and effort statistics by Area for each salmon species.

3.3.4.1 Nitinat Chum Salmon Management Strategy

Of importance to the Pacific Salmon Treaty is the WCVI group of Chum salmon populations returning to the Nitinat watershed (Nitinat River and Nitinat Lake and tributaries). This Nitinat stock is the most significant WCVI group of populations and includes production from a major CDFO hatchery on the Nitinat River.

The Nitinat commercial Chum salmon fishery (Area 21), along with the ESSR fisheries (Area 22), is the largest fishery on the WCVI and targets returning Nitinat River hatchery Chum salmon. In the past, catch of non-WCVI Chum salmon has occurred in the outside seaward portion of the Nitinat fishing area. Management actions have been implemented to reduce this catch of non-WCVI Chum salmon by restricting the Nitinat fishery to subareas closer to the terminal area (i.e. outlet of Nitinat Lake). More recently, First Nations' ESSR harvests have occurred within Nitinat Lake (Area 22). Historical Chum salmon harvest is shown in Table 3-10.

The spawning escapement target for Chum salmon returning to the Nitinat system (i.e. Nitinat River and other tributaries to Nitinat Lake) has been set at 175,000 fish (CDFO 2013). The target distribution of this spawning escapement between Nitinat River and the tributaries to Nitinat Lake has varied slightly over the last number of years; currently it is set at 125,000 into the Nitinat River and 50,000 into other Nitinat Lake tributaries. This spawning escapement requirement is set at a level to ensure that all run timing components are represented. Additional allocation for First Nations FSC requirements and hatchery requirements (i.e. broodstock) may total up to 50,000 Chum salmon. Consequently, the *gross escapement requirement* (spawning escapement + FN FSC requirements + hatchery requirements) for this Nitinat group of populations is 225,000 Chum salmon (CDFO 2013). Historical Chum salmon escapements are shown in Table 3-10.

A pre-season forecast of current year returns for the Nitinat stock is approximated based on broodyear escapements, estimated freshwater production from natural spawners, hatchery production, average age at return, environmental and ocean condition factors during brood ocean entry and estimated marine survival of each broodyear class.

A pre-season fishing plan is commonly developed by the end of the summer and is based on the forecast of current year returns. The fishery is managed through a fixed-harvest rate strategy to limit the harvest rate to approximately 25%. The allowable effort of the commercial fleets was determined from an analysis of the average exploitation rate per unit effort in Nitinat Chum salmon fisheries over the last decade.

In addition to concerns over catch of non-WCVI Chum salmon, early season opportunities are constrained by concerns over bycatch of Interior Fraser River Steelhead (Thompson River) and passing stocks of Coho salmon. Since 1995, these bycatch concerns in the commercial Nitinat fishery have been addressed by:

- *delayed opening dates*: the first commercial gill net fishery, if conducted, has been delayed until the beginning of October since 1998;
- *reduced fishing areas*: the traditional fishing area has been reduced to within a one-mile boundary between lines true south from Pachena and Dare Points (a portion of Area 21);
- *increased use of weed lines*: a 1.2m - 2.0m weed line for gill nets; and,
- *species selective fishing techniques*: non-retention of Coho salmon and Steelhead.

Additionally, to reduce bycatch mortality and to improve the quality of catch data, the following measures have been implemented in the Nitinat fisheries:

- *daylight fishing only*: gill net fishery;

- *mandatory functional revival tanks*: in purse seine and gill net fisheries;
- *on-board observers*: portions of purse seine and gill net fleet; and,
- *logbooks and weekly hail-ins*: purse seine and gill net fisheries.

The in-season management of the Nitinat fisheries (Area 21 and 22) are based on achieving the *gross escapement requirement* into the Nitinat watershed. Broodstock collection success, hatchery swim-ins, and visual surveys of Nitinat River spawners are used to determine whether the weekly *escapement milestone levels* into Nitinat Lake are achieved (CDFO 2013). Weekly fisheries are generally scheduled in Area 21 and surrounding waters to harvest any identified surplus.

First Nations FSC fisheries remain a priority and occur in terminal areas of Nitinat (Areas 21 and 22). First Nations ESSR fisheries can also occur in Nitinat Lake (Area 22) targeting returns surplus to spawning and hatchery requirements.

3.4 Annual Fishery Descriptions

Fisheries targeting Chum salmon in Johnstone Strait, Strait of Georgia, Fraser River, and on the West Coast of Vancouver Island (WCVI) generally begin in late September and end in November. These fisheries are directed at fall Chum salmon.

In addition to these directed fisheries, a bycatch of Chum salmon may occur in fisheries directed at Sockeye and Pink salmon earlier in the season. This Chum salmon bycatch is assumed to be comprised mainly of summer Chum salmon. The summer Chum salmon harvest in boundary area fisheries are presented in Table 3-4.

The main components of the Inside South Coast (ISC) Chum salmon return was expected to be both Fraser and non-Fraser stocks. These stocks are typically dominated by four year old fish from a below average 2009 brood return which out-migrated to the ocean in 2010. It was quite apparent that other salmon species that also out-migrated in 2010 encountered lower productivity and reduced survivals (Pink and Coho salmon returns in 2011). The pre-season expectation for ISC Chum salmon suggested low to near target returns to the area.

Table 3-4. Estimated harvest of summer Chum salmon by PFMA for commercial, research and test fishing vessels 1 (July through the second week of September), 2004–2013.

Year	Statistical Weeks	Pacific Fishery Management Area					Total
		18	19	20	21	29 ²	
2004	7/1–9/2	0	0	17	0	686	703
2005	7/1–9/2	0	0	0	0	115	115
2006	7/1–9/2	0	0	52	0	419	471
2007	7/1–9/2	0	0	0	0	49	49
2008	7/1–9/2	0	0	0	0	67	67
2009	7/1–9/2	0	0	0	0	27	27
2010	7/1–9/2	0	0	16	0	384	400
2011	7/1–9/2	0	0	0	0	883	883
2012	7/1–9/2	0	0	0	0	125	125
2013	7/1–9/2	0	0	0	0	27	27
Average		0	0	9	0	278	287

(1) Does not include PSC test fisheries harvest.

(2) Majority of this harvest is fresh water based and likely of Fraser River origin.

The following sections describe the fisheries directed at fall Chum salmon in each of the Southern B.C. areas in 2013.

3.4.1 Johnstone Strait

The Johnstone Strait test-fishery provided timing and abundance information for the 2013 return, which are important in assessing the performance of the 20% fixed exploitation rate strategy. It also provided an index of abundance, used to determine the likelihood of the number of returning Chum salmon being over the 1.0 million critical threshold level (requirement for commercial openings). Catch per unit effort in the test fishery was relatively strong and it was determined that the returning abundance was likely higher than the critical threshold required for continuation of commercial fisheries. All subsequent commercial openings in Johnstone Strait were prosecuted as planned. Age composition derived from the test-fishery and commercial samples was dominated by 4 year olds throughout the season. Escapements and catches in 2013 suggested returns were average to below average in many ISC Chum. Table 3-5 outlines the duration and Chum salmon harvest, of fishery openings during the 2013.

In 2013, there were two competitive commercial purse seine openings; three gill net openings in and one troll individual transferable quota (ITQ) fishery in Johnstone Straits (Area 12 and Area 13). The total 2013 commercial harvest for Johnstone Strait, including test fishery harvest, was 635,813 Chum salmon (Table 3-5). There were no directed commercial Chum salmon fisheries in the terminal areas of Johnstone Strait. Recreational harvest was low in Johnstone Strait (estimated at <200 Chum salmon; Table 3-5) with little monitoring during the month of October. First Nations FSC harvest was estimated at 13,444 in the Johnstone Strait area (Table 3-5). The total estimated harvest of Chum salmon in Johnstone Strait in 2013 was 649,400.

Table 3-5. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in Johnstone Strait by PFMA, 2013.

Gear Type	Area	Fishery Dates		Fishery Duration (hours)	Estimated Harvest
		Start	End		
Purse Seine (Test)	12	Sep. 16	Oct. 29	n/a	38,426
Purse Seine	12	Sockeye/Pink Directed ¹			7,538
	13				731
	12	Oct. 2	Oct. 3	16	105,028
	13	Oct. 2	Oct. 3	16	144,478
	12	Oct. 21	Oct. 21	10	64,992
	13	Oct. 21	Oct. 21	10	109,659
Gill Net	12	Oct. 7	Oct. 9	41	31,350
	13	Oct. 7	Oct. 9	41	14,422
	12	Oct. 15	Oct. 17	41	33,402
	13	Oct. 15	Oct. 17	41	12,411
	12	Oct. 25	Oct. 27	41	14,968
	13	Oct. 25	Oct. 27	41	16,034
Troll ²	12	Sep. 27	Oct. 13	n/a ²	0
	13	Sep. 27	Oct. 13	n/a ²	33,415
	12	Oct. 15	Nov. 3	n/a ²	0
	13	Oct. 15	Nov. 3	n/a ²	8,959
Recreational ³	12	n/a	n/a	n/a	41
	13	n/a	n/a	n/a	102
First Nations ³	12	n/a	n/a	n/a	3,565
	13	n/a	n/a	n/a	9,879
Total					649,400

(1) This is bycatch from Fraser Sockeye and Pink salmon directed fisheries that took place during early September.

(2) The troll fishery was opened from Sep. 27 to Nov. 3. The fishery was an effort based quota over that time period. Statweeks 9/4 to 10/2 are grouped and weeks 10/3 to 10/5 are grouped.

(3) The recreational and First Nation fisheries time periods varied over the season.

3.4.2 Strait of Georgia

In 2013, the total return of Chum salmon to Area 14 systems was within the preseason forecasted range (198,000 and 297,000). Commercial openings targeted Chum salmon returning to the three major systems in this area (Puntledge, Little Qualicum and Big Qualicum rivers). There were two competitive gill net openings and one purse seine opening in Area 14. Total commercial catch associated with Area 14 fisheries was 30,883 Chum salmon (Table 3-6).

In-season assessments in Area 18 did not identify sufficient escapement levels in the Cowichan River to initiate fisheries. Test fishery harvest was a total of 8,270 Chum salmon (Table 3-6).

The recreational creel survey within the Strait of Georgia area was not conducted after September in 2013.

The harvest by First Nations in the Strait of Georgia in 2013 was estimated to be 4,315 Chum salmon (Table 3-6).

The total estimated harvest of Chum salmon in the Strait of Georgia in 2012 was 53,461.

Table 3-6. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in the Strait of Georgia by PFMA, 2013.

Gear Type	Area	Fishery Dates		Fishery Duration (hours)	Estimated Harvest
		Start	End		
Purse Seine (Test)	18-19	Oct. 28	Dec. 3	n/a	8,270
Purse Seine	14	Nov. 6	Nov. 14	210	0
Gillnet	14	Nov. 3	Nov. 5	50	18,531
	14	Nov. 6	Nov. 20	354	52,352
	17	Nov. 3	Nov. 4	24	9,993
Recreational ¹	14-19, 28, 29 ³	May 1	Sep. 30	n/a	0
First Nations ²	14	n/a	n/a	n/a	30
	15	n/a	n/a	n/a	150
	16	n/a	n/a	n/a	0
	17	n/a	n/a	n/a	0
	18	n/a	n/a	n/a	3,030
	19	n/a	n/a	n/a	1,105
	28	n/a	n/a	n/a	0
Total					53,461

(1) The recreational fishery was not monitored after September; harvest of Chum salmon was likely very low.

(2) The First Nation fisheries time periods varied over the season.

(3) Marine subareas of Area 29, i.e. does not include an estimate for Chum salmon harvest in the tidal portion of the lower Fraser River downstream of Mission BC (Appendix B1).

(n/a) not applicable.

3.4.3 Fraser River

Directed Chum salmon fisheries occur within the Fraser River for all fishing sectors, including First Nations, recreational, commercial, and test fisheries. In recent years, significant conservation measures have been implemented in-river during the Fraser River Chum salmon migration period in order to protect co-migrating stocks of concern, particularly Interior Fraser River Coho salmon and Interior Fraser River Steelhead. Depending on the fishery, these measures have included time and area closures, gear restrictions and release requirements.

The escapement objective for Fraser River Chum salmon is 800,000 spawners. As described in section 3.3.3 (*Fraser River Chum Salmon Management Strategy*), Chum salmon fishing opportunities are provided based on in-season assessments of run-size, using CPUE information from the CDFO Albion Chum salmon test fishery. The initial in-season estimate of terminal abundance provided to the US on October 16, 2013 was 1,489 million Chum salmon. A subsequent in-season estimate of 1,634 million was provided on October 21.

Table 3-7. Estimated harvest of Chum salmon from salmon fisheries occurring in Area 29 (both marine and in-river) and in Region 2 (non-tidal) of the Fraser River, 2013.

Fishing Sector	Fishery Description	Target Species	Fishery Dates		Fishery Duration	Assessment Period ¹		Estimated Harvest
			Start	End		Start	End	
Total	Adriatic Chinook	Chinook	25-Apr	20-Oct	176 days	n/a	n/a	1,000
	Adriatic Chum	Chum	1-Sep	25-Nov	56 days	n/a	n/a	8,500
	Wharmouth Sechelt	Sechelt	24-Jul	4-Oct	107 days	n/a	n/a	1,100
	Commemorial Sechelt	Sechelt	16-Jul	16-Sep	62 days	n/a	n/a	0
Area 29 Subtotal								10,600
Commercial ²	Area 2 Gill Net	Chum	24-Oct	24-Oct	1 day	n/a	n/a	98,012
	Area 2 Seine	Pink	1-Sep	18-Sep	17 days	n/a	n/a	505
		Chum	17-Oct	17-Oct	1 day	n/a	n/a	6,100
Commercial Subtotal								104,617
Recreational ^{3,4,5}	Fraser River Mariculture	Mixed	27-Jul	15-Aug	20 days	27-Jul	15-Aug	0
		Mixed	16-Aug	16-Oct	125 days	16-Aug	16-Sep	15
	Chilliwack River	Coho	1-Jul	16-Sep	46 days	15-Sep	15-Nov	2,400
		Mixed	1-Jul	16-Oct	105 days			
	Sturgeon Slough	Mixed	1-Jul	16-Oct	97 days	12-Oct	16-Nov	1,000
Recreational Subtotal								3,415
First Nations ^{6,7}	FSC	Mixed	11-May	6-Sep	n/a	n/a	n/a	200
		Chum	1-Oct	26-Nov	n/a	n/a	n/a	27,216
	ESRD	Chum	n/a	n/a	n/a	n/a	n/a	45,176
	EDS	Pink	16-Sep	26-Sep	n/a	n/a	n/a	1,500
		Chum	16-Oct	6-Nov	n/a	n/a	n/a	106,018
First Nations Subtotal								154,910

(1) Assessment Period Start and End dates refer to the period of assessment for the recreational fisheries only.

(2) Fishery Start and End Dates for the commercial and First Nations fisheries reflect the first and last date the fisheries were permitted; the fisheries were not continuous between these dates.

(3) Fishery Start and End Dates for the recreational fisheries reflect the period the fishery was opened to the retention of Chum salmon; retention of other salmon species may have been allowed beyond these dates.

(4) The recreational estimates are for the periods of assessment; assessment periods may or may not coincide with Chum retention fishery opening and closing dates.

(5) Recreational estimates are preliminary.

(6) The Fishery Duration represents the maximum number of days provided in First Nations individual First Nations may have fished for shorter periods.

(7) The Fraser River maximum recreational fishery was closed to salmon fishing from Aug 16 to Aug 16, 2013.

n/a: not applicable

First Nations FSC gill net fisheries targeting Chum salmon were initiated October 1 following a closure period to protect co-migrating Interior Fraser Coho salmon. These fisheries harvested a total of 27,216 Chum salmon. First Nations Economic Opportunity (EO) fisheries directed at Chum salmon in 2013 harvested 106,018 Chum (Table 3-7). In addition to these fisheries, certain

First Nations groups were also provided access to Chum salmon that returned to hatchery facilities but that were not required for broodstock. A total of 40,176 Chum salmon deemed to be surplus to spawning requirements (ENR) were provided to First Nations from various hatchery facilities in 2013.

First Nations FNE (Chum) and Hatchery-derived Salmon accounting in 2013 had received harvest of Fraser Chum salmon, harvesting 219 Chum salmon (Table 3-7). An additional 1,940 Chum were taken as bycatch in Hatchery-derived Economic Opportunity Fisheries in 2013.

The total harvest of Chum salmon in all Fraser River First Nations Fisheries (including Hatchery ENR fish) was 188,370 in 2013 (Table 3-7 and Table 3-8).

Table 3-8: Estimated harvest of Chum salmon from First Nations and commercial salmon fisheries occurring in June 29 and Fraser River, 2004-2013

Year	First Nations			Commercial			Total
	FNE	ENR	ENR	Harvest ENR June 12-18	Other and FNE Total Fisheries	Other Commercial Fisheries	
2004	15,400	45,400	46,400	55,400	15,400	0	116,600
2005	7,400	45,400	46,400	55,400	15,400	0	111,700
2006	15,400	116,700	46,400	116,700	15,400	0	394,200
2007	15,400	27,400	27,400	46,400	15,400	0	121,700
2008	15,400	46,400	46,400	46,400	15,400	0	170,600
2009	15,400	46,400	46,400	46,400	15,400	0	169,600
2010	15,400	0	15,400	0	15,400	0	30,800
2011	15,400	46,400	46,400	46,400	15,400	0	169,600
2012	15,400	116,700	46,400	116,700	15,400	0	394,200
2013	27,400	127,400	46,400	116,700	15,400	0	329,300
Average	15,400	47,400	46,400	66,400	15,400	0	287,000

(1) Includes "experimental" fisheries taking surplus harvest under the CDFP Fisheries Training Program; none are applicable.

The major Fraser River watershed recreational salmon fisheries harvesting Chum salmon in 2013 include significant fisheries occurring in the lower Fraser River mainstem, the Chilliwack River and the Skagit River, and minor salmon fisheries occurring in the Harrison River and the Nicomen Slough-Narada Creek System. The lower four systems are tributaries in the Fraser River in the lower Fraser Valley. The recreational fisheries occurring in the Harrison and Skagit rivers were not assessed in 2013.

The lower Fraser River mainstem was open to recreational angling and the harvest of Chum salmon on two separate occasions in 2013. The initial opening occurred from July 27 to August 15, 2013 (daily retention limit of 2) and was followed by a closed period of approximately 5 weeks. The lower Fraser River mainstem was subsequently opened from August 15 to December 15, 2013 (daily retention limit of 2). In 2013, the mainstem recreational fishery was assessed from July 27 to August 15 and August 15 to September 30, 2013, estimates of 77 and 155 Chum salmon were harvested and released, respectively (Table 3-9). The recreational harvest in 2013 was notably less than in 2012 (Table 3-9). However, assessment in 2013 was shortened

was recorded earlier than in 2012. In 2012, approximately 50% of the reported harvest occurred in the October 1 to November 30 period.

The Chittawab Creek recreational fishery was open to the collection of *C. bairdi* offshoot from July 1 to November 30, 2012 with a season limit of 1. This fishery was opened from September 15 to November 15, 2013, however, of 1,400 and 12,000 *C. bairdi* offshoot were harvested and released, respectively (Figure 1-7). Working in the House River, numerous recreational fishers, the estimated *C. bairdi* offshoot catch in 2012 decreased from 2011 due to the Chittawab Creek recreational fishery (Figure 1-7). However, within the House fishery, the fishery management was identical, consistent between 2012 and 2013.

The Hartman River, House River and Neenah-Booth recreational fisheries were open to the collection of *C. bairdi* offshoot with a season limit of 1. The Neenah-Booth fishery was opened from October 1 to November 30, 2012. Estimates of 1,400 and 1,200 *C. bairdi* offshoot were harvested and released, respectively, for the fishery (Figure 1-7) and were a significant increase over the estimates reported for 2011 (Figure 1-7). Working in the Chittawab Creek recreational fishery, the fishery management in 2012 was comparable to the management in 2011. Recreational fisheries occurring in the Hartman and House rivers were not assessed in 2012.

In total, the assessed recreational fisheries occurring in the House House River contributed in 2012 estimates of 1,400 and 12,000 *C. bairdi* offshoot were harvested and released, respectively (Figure 1-7 and Figure 1-7).

Commercial fisheries in the House House River (i.e., House B gill net and net-potting grounds) were open September 1 to mid-November during the House House *C. bairdi* regulation period. In recent years, the House has been assessed over two October to provide additional protection to House House headland. As a result, commercial House B gill net *C. bairdi* offshoot fisheries were typically open during the last week of October and first week of November. In 2012, there was one House B commercial opening in the House River on October 19 that harvested over 10,000 *C. bairdi* offshoot. The House B crew that also conducted a limited participation *C. bairdi* offshoot harvest fishery off the mouth of the House River on October 17 that harvested 1,200 *C. bairdi* offshoot (Figure 1-7). Finally, there is also had a demonstration fishery, against a House River flag offshoot off the mouth of the House River. This fishery harvested the *C. bairdi* offshoot between September 1 and September 10.

Copyright © 2006 John Wiley & Sons, Inc. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, except as may be permitted in writing by John Wiley & Sons, Inc.

Year	Government Expenditure (Million US\$)					
	Health	Education	Defense	Other	Total	Share of GDP
1990	100	150	200	50	500	1.1%
1991	105	155	205	55	520	1.1%
1992	110	160	210	60	540	1.1%
1993	115	165	215	65	560	1.1%
1994	120	170	220	70	580	1.1%
1995	125	175	225	75	600	1.1%
1996	130	180	230	80	620	1.1%
1997	135	185	235	85	640	1.1%
1998	140	190	240	90	660	1.1%
1999	145	195	245	95	680	1.1%
2000	150	200	250	100	700	1.1%
2001	155	205	255	105	720	1.1%
2002	160	210	260	110	740	1.1%
2003	165	215	265	115	760	1.1%
2004	170	220	270	120	780	1.1%
2005	175	225	275	125	800	1.1%
2006	180	230	280	130	820	1.1%
2007	185	235	285	135	840	1.1%
2008	190	240	290	140	860	1.1%
2009	195	245	295	145	880	1.1%
2010	200	250	300	150	900	1.1%
2011	205	255	305	155	920	1.1%
2012	210	260	310	160	940	1.1%
2013	215	265	315	165	960	1.1%
2014	220	270	320	170	980	1.1%
2015	225	275	325	175	1000	1.1%
2016	230	280	330	180	1020	1.1%
2017	235	285	335	185	1040	1.1%
2018	240	290	340	190	1060	1.1%
2019	245	295	345	195	1080	1.1%
2020	250	300	350	200	1100	1.1%

© 2011, a group of 111,000 Hong Kong citizens, with the support of a consortium including Microsoft, Apple, Google, Facebook, YouTube, and the Internet Archive, to the Hong Kong Police and the central government, all the Chinese Mainland people, 111,000 people, 111,000 people.

[illegible][illegible][illegible]

© 2006 by the author(s). All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without permission in writing from the publisher.

© 2004 Blackwell Publishing Ltd, *Journal of Internal Medicine* 255: 105–112

© 2006 The Authors
Journal compilation © 2006 Blackwell Publishing Ltd

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

1. The first step is to identify the problem. In this case, the problem is that the company is not meeting its sales targets.

© 2006 The Authors
Journal compilation © 2006 Blackwell Publishing Ltd

© 2006 The Authors
Journal compilation © 2006 Blackwell Publishing Ltd

and Puntledge river Hatcheries. There is one enhancement facility in the Burrard Inlet (Capilano River Hatchery: limited Chum salmon hatchery production), and the Squamish watershed (Tandusht Creek Hatchery) and four in the Fraser River watershed (Inch Creek, Chilliwack River, and Chehalis River Hatcheries and Wenner Creek Spawning Channel). Enhancement on the Capilano River is opportunistic, only using Chum salmon that return to the hatchery facility for broodstock.

3.5.2 West Coast Vancouver Island

In 2013, Nitinat Chum salmon spawning escapement was estimated at 25,000 (Table 3-10) which is a significant reduction compared to the 2012 return, and the lowest since 2000. In other WCVI Areas, Chum salmon spawning escapements were poor and below average in all areas except Area 26 (Table 3-12). Spawning escapement estimates given for Areas 23-26 in Table 3-12 are derived from index system assessments in each Area. From historical surveys of Chum salmon streams in each Area, the proportion each stream contributed (on average) to the total spawning escapement for the Area is estimated. Spawning escapement estimates from index systems currently surveyed annually, are expanded to an Area spawning escapement to account for those streams not surveyed.

Table 3-12 WCVI Chum salmon spawning escapement estimates¹ by PFMA, 2004-2013

Year	WCVI Pacific Fishery Management Area			
	Area 23 (Barkley)	Area 24 (Clayoquot)	Area 25 ² (Nootka)	Area 26 (Kyuquot)
2004	115,452	111,700	241,870	76,708
2005	47,546	55,047	114,007	96,709
2006	23,720	117,407	41,278	78,479
2007	86,012	42,272	45,178	91,087
2008	58,070	25,878	46,497	55,342
2009	27,020	25,302	42,178	52,852
2010	58,752	44,709	71,009	76,628
2011	120,092	149,164	61,989	125,989
2012	40,439	42,276	49,144	49,923
2013	25,302	44,178	69,877	74,412
Average	74,740	72,308	47,071	66,154

(1) Index system spawning escapements expanded to total spawning escapement for the Area

(2) Nootka spawning escapements include hatchery river systems

4 United States Chum salmon

4.1 Washington Run Sizes, Catches, and Spawning Escapements

Tables 4-1 through 4-3 provide pre-season forecasts of run size, post-season estimates of run size and post-season estimates of spawning escapement for the Chum salmon runs returning to Puget Sound and coastal Washington areas. The tables present estimates for three major Chum salmon groupings, defined by their return timings.

Summer Chum Salmon

The Strait of Juan de Fuca summer Chum salmon post-season run size of 14,801 was 2.2 times larger than the pre-season forecast and was 2.2 times larger than the 2004 – 2012 average post season run size. The Hood Canal summer Chum salmon run size of 24,325 was 23% larger than the pre-season forecast and was 97% of the 2004 – 2012 average post-season run size. Approximately 95% of the combined Strait of Juan de Fuca and Hood Canal summer Chum salmon run went to escapement. The summer Chum salmon pre-season forecast for South Puget Sound for 2013 was estimated at approximately 53,500 fish, which is 88% of the average return for 2004-2012. The post-season run size estimate of South Puget Sound summer Chum salmon in 2013 was 25,317 fish, or 43% of the pre-season forecast, and was 42% of the 2004-2012 average post-season run size.

Table 4-1. Puget Sound summer Chum salmon pre-and post-season estimates of run size and spawning escapements, 2004–2013.

Region	Type	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Strait of Juan de Fuca	Pre-Season	4,729	6,803	8,238	8,566	5,969	5,198	3,991	5,308	5,915	6,603
	Post-Season	9,361	9,730	8,280	3,324	3,576	5,147	9,332	5,705	6,337	14,801
	Escapement	9,280	9,619	8,181	3,219	3,449	5,029	9,179	5,675	6,304	14,727
Hood Canal	Pre-Season	18,078	18,060	19,780	23,729	20,159	18,009	5,998	9,050	8,970	19,798
	Post-Season	88,236	16,313	29,990	12,838	17,618	9,200	12,957	7,170	31,134	24,325
	Escapement	69,565	15,311	26,418	10,539	15,112	7,236	12,533	6,914	29,855	22,618
South Puget Sound	Pre-Season	99,317	38,334	55,300	39,840	64,229	57,352	62,991	62,623	41,889	53,492
	Post-Season	178,199	44,993	78,797	57,786	32,065	25,030	49,098	39,823	38,936	25,317
	Escapement	120,782	24,701	63,345	52,669	20,872	21,639	65,808	38,517	24,522	20,593




Figure 4-1. Puget Sound summer Chum salmon abundance, 2004-2013, by region of origin (JDF: Strait of Juan de Fuca, HOOD: Hood Canal, SPS: South Puget Sound).

Fall Chum Salmon

The Puget Sound fall Chum salmon post-season assessment total was 2,141,514 fish, or 271% of the preseason forecast and was approximately 340% of the 2004-2013 average run size. Regional post-season numbers varied in their deviation from the forecasted numbers, ranging from 441% (Hood Canal) to 64% (Skagit River) of forecasted values (Table 4-2, Figure 4-2). The Washington coastal (Willapa Bay and Grays Harbor) fall Chum salmon post-season assessment total was 69,088 fish, or approximately 104% of the preseason forecast and was 104% of the 2004-2012 average run size.




Figure 4-2. Puget Sound fall Chum salmon abundance, 2004-2013, by region of origin (JDF: Strait of Juan de Fuca, NOOK: Nooksack-Samish, SKAG: Skagit, STLSNO: Stillaguamish/Snohomish, SPS: South Puget Sound, HOOD: Hood Canal).

Winter Chum Salmon

The winter Chum salmon 2013 post-season run size assessment for South Puget Sound was 34,839, or approximately 57% of the preseason forecast and was 41% of the 2004-2012 average run size (Table 4-3, Figure 4-3).




Figure 4-3. South Puget Sound winter Chum salmon abundance, 2004-2013

Table 4-2. Washington fall Chum salmon pre-and post-season estimates of run size and spawning escapements, 2004–2013.

Region	Type	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Willapa Bay	Pre-Season	79,183	63,441	38,754	23,798	40,022	7,139	33,442	22,254	28,273	35,584
	Post-Season	114,758	28,131	23,486	17,668	12,989	17,444	26,701	69,802	43,069	27,564
	Escapement	84,926	11,983	15,250	17,392	9,511	12,743	25,204	66,871	26,640	24,979
Grays Harbor	Pre-Season	45,352	38,038	29,370	12,000	9,903	17,665	18,492	33,669	24,603	30,716
	Post-Season	34,274	20,596	15,786	12,136	8,879	19,776	43,582	50,494	40,561	41,524
	Escapement	17,494	13,483	11,268	11,342	6,249	15,216	34,644	30,101	27,876	22,519
Strait of Juan de Fuca	Pre-Season	2,438	3,460	3,830	2,143	2,745	2,587	2,222	1,618	1,203	813
	Post-Season	3,233	2,382	1,567	769	841	1,462	1,849	4,839	3,174	1,468
	Escapement	2,739	2,034	1,313	503	454	1,305	1,343	4,707	2,973	1,394
Nooksack/Samish	Pre-Season	78,484	126,869	135,100	19,414	130,070	69,145	61,827	31,924	34,739	52,585
	Post-Season	89,901	67,657	66,610	48,631	27,471	32,007	48,895	152,166	39,080	68,170
	Escapement	53,563	47,644	29,289	21,571	16,257	26,267	30,509	77,995	21,684	48,500
Skagit	Pre-Season	109,715	25,695	164,094	90,481	132,036	26,828	50,226	26,834	59,167	15,325
	Post-Season	171,200	53,672	213,490	37,627	29,946	29,479	46,149	15,851	41,600	9,786
	Escapement	150,196	34,600	105,239	19,576	22,552	26,744	39,911	15,494	36,601	8,554
Stillaguamish/Snohomish	Pre-Season	264,542	225,113	445,800	287,993	229,251	92,477	90,660	71,277	86,598	48,884
	Post-Season	382,872	84,791	479,987	235,799	125,064	37,689	109,358	61,580	40,002	35,144
	Escapement	212,463	38,788	272,925	43,664	38,553	29,378	46,628	60,792	35,450	30,317
South Puget Sound	Pre-Season	470,048	655,742	466,700	408,040	686,511	348,333	650,986	487,514	323,928	349,623
	Post-Season	1,022,749	414,939	939,638	1,036,774	464,193	344,854	623,197	545,192	613,564	599,668
	Escapement	356,712	137,442	369,035	369,455	152,579	150,887	201,992	175,989	191,290	186,908
Hood Canal	Pre-Season	501,100	749,593	668,400	587,155	668,397	374,929	477,409	352,019	426,675	323,597
	Post-Season	1,173,077	356,243	745,675	676,602	553,122	446,476	416,461	611,349	684,662	1,427,277
	Escapement	229,893	100,242	185,875	162,425	72,012	55,879	49,484	103,383	91,973	240,111

Table 4-3. Washington winter Chum salmon pre- and post-season estimates of run size and spawning escapements, 2004–2013.

Region	Type	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
South Puget Sound	Pre-Season	76,464	142,406	149,200	81,065	98,922	83,380	89,293	55,923	51,599	61,113
	Post-Season	134,002	72,730	75,833	105,933	89,271	35,544	102,235	121,079	25,522	34,839
	Escapement	98,580	43,908	58,786	73,442	45,162	15,486	79,550	74,110	18,987	21,754

Paragraph 3 of the Chapter 6, Annex IV of the Treaty requires that Canada and the United States assess catch levels of summer Chum salmon caught during the August 1 through September 15 time period in boundary area fisheries. Table 4-4 provides Chum salmon harvest during this summer Chum salmon accounting period for U.S. boundary area fisheries. Table 4-5 provides Chum salmon harvest information from the Strait of Juan de Fuca (SJF) fisheries in Areas 4B, 5, and 6C and the San Juan Islands/Point Roberts (SJI/PR) fisheries in Areas 7 and 7A (Appendix B1). Table 4-5 also includes annual Chum salmon harvest totals for the Puget Sound and the Washington Coastal areas.

Table 4-4. Harvest of Chum salmon in the Strait of Juan de Fuca (SJF) and the San Juan Islands (SJI) commercial fisheries during the summer Chum salmon accounting period, July 1–September 15, 2004–2013.

Year	Time Period ¹					Yearly Total
	7/1–8/18	8/19–8/25	8/26–9/1	9/2–9/8	9/9–9/15	
SJF/HC GSI ₂	0.68	0.40	0.45	0.14	0.07	
2004	162	0	15	31	25	233
2005	29	17	17	3	16	82
2006	98	58	10	8	3	177
2007	39	0	0	0	9	48
2008	240	1	0	0	29	270
2009	13	0	0	17	52	82
2010	66	140	46	39	6	297
2011	31	0	3	27	20	81
2012	135	0	0	0	0	135
2013	52	21	115	129	462	779
Averages	87	24	21	25	62	218

(1) Indicates cumulative catch of Chum salmon through this period (GSI would be applied to values to get numbers of HC/SJF summer Chum salmon). Time period notation is month/day (e.g. 7/1–8/11 refers to July 1 to August 11).

(2) Estimated proportions of Hood Canal/Juan de Fuca summer Chum salmon in Boundary area waters (GSI from 1995–1997).

Table 4-5. Harvest of summer, fall, and winter Chin salmon in SIF, SJUPR, Puget Sound and Washington coastal areas, 2004-2013¹

Year	Region			
	SIF (Areas 4B, 5, 6C)	SJUPR (Areas 7, 7A)	Puget Sound ²	WA Coast ³
2004	4926	106,441	2,076,012	48,049
2005	1,771	77,096	608,496	23,870
2006	4,483	106,614	1,508,976	13,932
2007	6,705	27,397	1,444,200	2,728
2008	5,695	73,374	964,076	7,067
2009	746	24,244	637,481	10,351
2010	1941	23,652	949,870	12,903
2011	1,930	70,363	1,009,490	28,251
2012	637	73,235	1,083,978	30,007
2013	1242	80,509	1,067,244	23,060
Averages	3,008	72,548	1,206,500	20,019

(1) Includes recreational harvest. 2013 recreational harvest estimates are preliminary.

(2) All other Puget Sound freshwater and marine harvest areas, except Strait of Juan de Fuca or San Juan Island/Point Roberts Fisheries.

(3) WA Coast combines harvest from Areas 1-4 including Grays Harbor, Willapa Bay, Columbia River and Coos Bay.

4.2 United States Strait of Juan de Fuca Fisheries (Areas 4B, 5, 6C)

4.2.1 Management Intent

During the 2013 season, the management strategy for the Strait of Juan de Fuca (SJF) fishery consisted of maintaining reduced effort in this fishery and making management decisions based on the needs of Puget Sound stocks of Chum salmon. This fishery has experienced minimal effort in recent years, resulting in relatively low harvest levels over the period of 2004–2013 (Table 4-5). Harvest and effort for the period of 2004–2013 was well below levels observed in the late 1980s and 1990s.

4.2.2 Fishery Description

As in previous years, the fall Chum salmon fishery in Areas 4B, 5 and 6C was restricted to Treaty Indian fishers using gill nets. The Treaty Chum salmon fishery opened the week of October 6, 2013 with a schedule of six days per week and continued through November 8, 2013. A total of 1,123 Chum salmon was harvested in this fishery, and there was a reported bycatch of 711 Coho salmon, 0 Steelhead, and 0 Chinook salmon.

In Areas 4B, 5, and 6C, the incidental harvests of Chum salmon prior to the Chum salmon-directed fishing season was relatively low in contrast to the recent ten year average. A total of 74 Chum salmon, including those harvested in SJF net fisheries, were recorded during the summer Chum salmon accounting period (July 1 – September 15). During the subsequent Coho salmon-directed fishery, an additional four Chum salmon were harvested.

The total 2013 Chum salmon catch by all gears (including recreational) in the SJF was 1,242 Chum salmon (Table 4-5).

4.3 San Juan Islands / Point Roberts Fisheries (Areas 7 and 7A)

4.3.1 Management Intent

Chum salmon fisheries in Areas 7 and 7A are regulated to comply with a base harvest ceiling of 130,000 Chum salmon, unless a critically low level of abundance ($< 1,000,000$) is identified for those populations migrating through Johnsons Strait (Inside Chum salmon). Chapter 5, Areas IV, specifies that Chum salmon-directed fishing is not allowed in Areas 7 and 7A before October 15. Paragraph 10 (a-b) specifies that for run sizes below the critical threshold, harvest of Chum salmon in Areas 7 and 7A will be limited to those taken incidentally to other species and in other minor fisheries, and shall not exceed 20,000.

4.3.2 Fishery Description

An estimated Fraser River Chum salmon run size of 1,000,000 was provided by Canada on October 15.

The U.S. Chum salmon fishery which had begun on October 10th continued through November 9th with weekly catch monitoring. The U.S. catch between October 10 and November 9th in Areas 7 and 7A was 79,350 Chum salmon. The Non-Treaty gill net and purse seine fleets were open daily October 12, 13, 15, and then open continuously from October 19 through November

66. The Treaty Indian gill net and purse seine fisheries were opened on October 15, and have run continuously from October 22 through November 16. Catch per vessel and effort were good throughout the fishery.

Seine Indian net and fisheries targeting offshore-matched Chin salmon were conducted from the end of Fraser River season (September 26), until September 30, with Chin salmon catches prohibited. From October 1 through November 6, net sets were open daily with Chin salmon catches allowed. Chin salmon catch in this fishery, between October 1 and October 15, was 16 fish. Effort was low and there was no net or fishing effort after early October.

There were 104 Chin salmon reported catches in Areas 7 and 14 during Fraser River approved fisheries and float salmon fisheries in August and early September. The total 104 Chin salmon catch by all gear (including recreational) in Areas 6, 7, and 14, reported through November 6, was 40,100. Catch distribution between Areas 7 and 14, was 72% and 28%, respectively. During the fall Chin salmon fisheries in Areas 6, 7, and 14, there was a reported harvest of 2 Chinook and 8,764 Chin salmon and 2 steelhead.

9 Stock Identification

9.1 Tagging of Adult Chin Salmon

No adult tagging projects were reported by either Canada or the United States in marine or nearshore areas in 2013.

9.2 Canada, Fin Marking and United River Tagging (CUT)

9.2.1 Canada

Thermal marking of adults (Volk *et al.* 1987) has been used to mark hatchery steelhead salmonids in B.C. since the early 1990's (Hargrave *et al.* 2001). The method involves manipulating the temperature of the marking water to at least 2°C in order to alter or destroy deposition chemistry of otolith layers. By systematically altering water temperatures over a period of time, banding patterns (i.e. a 'mark') unique to each hatchery can be created.

Thermal marking was first used on Chin salmon at Nahcotta Hatchery on the West Coast of Vancouver Island in 1995 and continues to be used as a means of identifying hatchery contribution to both fisheries and escapement. Thermal marking is currently the only mark being applied to Chin salmon from Nahcotta Hatchery (Table 5-1). Transplanting thermally-marked Chin salmon into Klamath River started with the 2004 brood and ended with the 2006 broodyear (Table 5-1).

Table 5.1. Estimates of C -test statistic with normally-matched weights from 500 C -test and C -weight function replications (1000 C -test)

Group Type	Source Function Between-Group		C-weight Function, Between-Group					
	Control Group of Factor	Reference Group	C-weight Group	C-weight Reference	Weight Group	Weight Reference	C-weight Group	Weight Reference
0000	1-1000-100	1-1000-100		100-100	100-100	100-100	100-100	
0001	10-100-100	1-1000-100		100-100	100-100	100-100	100-100	
0002	10-100-100	1-1000-100		100-100	100-100	100-100	100-100	
0003	1-1000-100			100-100	100-100	100-100	100-100	
0004	1-1000-100			100-100	100-100	100-100	100-100	
0005	1-1000-100			100-100	100-100	100-100	100-100	
0006	1-1000-100			100-100	100-100	100-100	100-100	
0007	1-1000-100			100-100	100-100	100-100	100-100	
0008	1-1000-100			100-100	100-100	100-100	100-100	
0009	1-1000-100			100-100	100-100	100-100	100-100	
0010	1-1000-100			100-100	100-100	100-100	100-100	
0011	1-1000-100			100-100	100-100	100-100	100-100	
0012	1-1000-100			100-100	100-100	100-100	100-100	
0013	1-1000-100			100-100	100-100	100-100	100-100	

A second function on the first C -test of function of which C -weight function, the group normally-matching weights for second replications of C -test statistic beginning with the 1000 bootstrap. The continued in 500 with normal weights of normally-matched C -test statistic to various size values of function group (Table 5.1). Starting a C -weight statistic, comparison of which sampling programs used in sampling it is comparing the normally-matched C -test statistic have occurred during the 100 C -test statistic values. Information from these programs has provided a better understanding of using and which functions for the C -test statistic population in function group.

[illegible]

Algorithm	Iterations	Min / Max Value	Average	Min / Max Variance	Final
My Solution (10)	100	1000	10.00	1.00	10.00
	200	1000	10.00	1.00	10.00
	300	1000	10.00	1.00	10.00
	400	1000	10.00	1.00	10.00
	500	1000	10.00	1.00	10.00
	600	1000	10.00	1.00	10.00
	700	1000	10.00	1.00	10.00
	800	1000	10.00	1.00	10.00
	900	1000	10.00	1.00	10.00
	1000	1000	10.00	1.00	10.00
Final Value	1000	1000	10.00	1.00	10.00

For all affine groups, we represent Γ as a set of matrices in $\mathrm{GL}(n, \mathbb{R})$ and represent the affine group action naturally as a mapping from $\mathbb{R}^n$ to $\mathbb{R}^n$. Affine groups acting on $\mathbb{R}^n$ can be thought of as n -dimensional transformations applied to the affine coordinates. We apply our theory to the well-known image registration problem of affine deformations $(\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b})$ (Fig. 1).

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–406

[illegible][illegible]

1.1.1. **General description of the project**

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

1.1.2. **General description of the project**

1.1.2.1. **General description**

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

Year	Project start	Project end	Project start	Project end
2010	2010	2010	2010	2010
2011	2011	2011	2011	2011
2012	2012	2012	2012	2012
2013	2013	2013	2013	2013
2014	2014	2014	2014	2014
2015	2015	2015	2015	2015
2016	2016	2016	2016	2016
2017	2017	2017	2017	2017
2018	2018	2018	2018	2018
2019	2019	2019	2019	2019
2020	2020	2020	2020	2020
2021	2021	2021	2021	2021
2022	2022	2022	2022	2022
2023	2023	2023	2023	2023
2024	2024	2024	2024	2024
2025	2025	2025	2025	2025
2026	2026	2026	2026	2026
2027	2027	2027	2027	2027
2028	2028	2028	2028	2028
2029	2029	2029	2029	2029
2030	2030	2030	2030	2030

1.1.2.2. **General description**

The project is a research project in the field of computer science, specifically in the area of artificial intelligence. The project is aimed at developing a system that can learn from data and make decisions based on that data. The project is funded by the European Union and the German government.

These ambiguous supplementation program fish were genotyped and identified by assigning them to populations in the Hood Canal summer Chum salmon genetic baseline. Genetic analysis was used to evaluate impacts of supplementation on the Hood Canal summer Chum salmon metapopulation (Small et al. 2009). WDFW also conducted juvenile stock assignment studies in the Strait of Juan de Fuca and Hood Canal during 2009 to 2013. The juvenile studies documented emigration timing and relative production of summer and fall Chum salmon in key populations. Further, 963 tissue samples were collected near Apple Cove Point in Puget Sound Area 9 in the eighth year of Chum salmon DNA collection from this test fishery.

Fisheries sampling in Area 7 and Area 7a was sampled and analyzed from the second of four years of Southern Endowment Funding project using microsatellites (Van Will et al. 2013).

Table 5-6: Chum salmon tissue samples collected in Areas 7 and 7a analyzed for GSI in 2012-2013

Year	Area 7	Area 7A
2012	1,526	429
2013	409	423

3.2.2 Baseline Collection for Genetic Stock Identification

3.2.2.2 Canada

In 2013, CEBCO continued to collect baseline GSI tissue samples of Southern British Columbia Chum salmon populations (Table 5-7).

Table 5-7: Fall Chum salmon GSI tissue collections from Southern B.C. by Conservation Unit, collection site, life stage, and influence¹ of enhancement, 2013

Conservation Unit	Collection Site	Life Stage	Enhancement Level ¹	Sample Size
Fraser Canyon	Stoke Cc	Adult	Natural	43
Fraser Canyon	Kayhawa Cc	Adult	Natural	47
George Strait	Obcevan	Adult	Natural	223
George Strait	Therabon R	Adult	Natural	144
Hove Sound/Burned Lake	Chickama R	Adult	Natural	146
Lang/Borough	Phillips R	Adult	Natural	57
Lower Fraser	Hasey Cc	Adult	Natural	66
Lower Fraser	Chilga Cc	Adult	Natural	1
Lower Fraser	Silverhops Cc	Adult	Natural	242
Lower Fraser	Hale Cc	Adult	Natural	147
Lower Fraser	Hopetide Slough	Adult	Natural	77
Lower Fraser	Wharton Cc	Adult	Natural	47
Lower Fraser	Bulford Cc	Adult	Natural	4

Conservation Unit	Collection Site	Life Stage	Enhancement Level ¹	Sample Size
Lower Fraser	Stuart Cr	Adult	Natural	27
Lower Fraser	Wharmock R	Adult	Natural	206
Lower Fraser	Wootz Cr	Adult	Natural	74
Lower Fraser	Big Silver Cr ²	Adult	Natural	1
Lower Fraser	Cashburn Cr ²	Adult	Natural	1
Lower Fraser	Saddle Cr ²	Adult	Natural	2
Lower Fraser	Wootz Cr ²	Adult	Natural	108
Lower Fraser	Porch Cr ²	Adult	Natural	163
Lower Fraser	Widgeon ²	Adult	Natural	135
Total				1969

(1) Enhancement level refers to the degree to which the CDFO Committee members believed hatchery production contributed to the returns to the collection site. Based partly on the enhancement history in the CDFO hatchery production database, enhancement level is subjectively classified as either *hatchery* (insignificant contribution of natural production to returns), *mixed* (both enhanced and natural production contribute significantly to returns) or *natural* (insignificant contribution of hatchery production to returns).

(2) indicates samples that have been collected but not analyzed for the microsatellite baseline.

3.3.2.1 United States

During 2013, WDFW and Washington Treaty Tribes continued baseline sampling of Washington State populations of Chum salmon (Table 5-8). Tissues were archived for future genotyping and will be added to the genetic baseline.

Table 5-8. Chum salmon GSI tissue collections from WA State, 2013

Collection Site / Run group / Locality	Enhancement Level ¹	Sample Size
Puyallup Hatchery - fall	hatchery	81
Cowwam Creek - fall	mixed	51
Sittigunomish River-fall	natural	236
Tulalip Hatchery - fall	hatchery	100
Snohomish River - fall	natural	106
Total		547

(1) Enhancement level refers to the degree to which the U.S. Committee members believed hatchery production contributed to the returns to the collection site: *hatchery* (insignificant contribution of natural production to returns), *mixed* (both enhanced and natural production contribute significantly to returns) or *natural* (insignificant contribution of hatchery production to returns).

REFERENCES CITED

- Beattie, W., G. Kirby, L. LeClair. 1996. The Stock Composition of 1995 Chum Salmon Fisheries in Washington: Strait of Juan de Fuca (Catch area 5) and Northern Puget Sound (Catch Areas 7&7a). Northwest Indian Fisheries Commission and WDFW Joint document.
- CDFO [Canadian Department of Fisheries and Oceans]. 2013. Pacific Region Integrated Fisheries Management Plan, Salmon, Southern B.C., June 1, 2013 to May 31, 2014. Fisheries Management Branch, Fisheries and Oceans Canada. Vancouver, British Columbia.
- Gazey, W.J. and R.V. Palermo. 2000. A preliminary review of a new model based on test fishing data analysis to measure abundance of returning Chum stocks to the Fraser River. DFO Canadian Stock Assessment Secretariat Research Document 2000/159. 30 p.
- Hargreaves, B., W. Luedke, and J. Till. 2001. Application of otolith thermal mass marking in British Columbia, Canada. NPAFC Technical Report No. 3. 2001 International Workshop on Salmonid Otolith Marking. March 21, 2001, Seattle, WA, U.S.A.
- Point No Point Treaty Tribes and Washington Department of Fish and Wildlife. 2014. Five-year review of the Summer Chum Salmon Conservation Initiative for the period 2005 through 2013: Supplemental Report No. 8, Summer Chum Salmon Conservation Initiative – An Implementation Plan to Recover Summer Chum in the Hood Canal and Strait of Juan de Fuca Region, September 2014. Wash. Dept. Fish and Wildlife. Olympia, WA. 237 pp., including Appendices.
- Small, M.P., K. Currens, T. H. Johnson, A. E. Frye and J. F. Von Bargaen 2009. Impacts of supplementation: Genetic diversity in supplemented and unsupplemented populations of summer Chum salmon (*Oncorhynchus keta*) in Puget Sound (Washington, USA). Canadian Journal of Fisheries and Aquatic Sciences 66(8):1216-1229.
- Van Will, P. J. R. Candy, and J. Zischke. 2013. Joint US and CA mixed-stock Chum fisheries sampling design and analysis. Report to Southern Endowment Fund: Project 57536.
- Volk, E.C., Schroder, S.L., and Fresh, K.L. 1987. Inducement of banding patterns on the otoliths of juvenile Chum salmon (*Oncorhynchus keta*). Proceedings of the 1987 northeast Pacific pink and Chum salmon workshop. Alaska Department of Fish and Game: 206-212.

APPENDICES

Appendix A. ANNEX IV, CHAPTER 6, OF THE PACIFIC SALMON TREATY

Southern British Columbia and Washington State Chum Salmon


The provisions of this Chapter shall apply for the period 2010 through 2018.

1. The Parties shall maintain a Joint Chum Technical Committee ("the Committee") reporting, unless otherwise agreed, to the Southern Panel and the Commission. The Committee will undertake to, *inter alia*:
 - a) maintain and present historical catch and escapement information for stocks relevant to the Treaty;
 - b) utilize available information to estimate and document stock composition and exploitation rates in fisheries of concern to the Treaty;
 - c) review annually the Parties' assessment of stock status and fisheries activities for Chum fisheries of concern to the Treaty;
 - d) identify high priority research and information needs for the Parties, including fishery and escapement monitoring and assessment, stock identification, and enhancement; and
 - e) periodically and/or when requested;
 - i. Exchange available information on the productivity and escapement requirements of stocks relevant to the treaty;
 - ii. Identify and document stocks of concern (with respect to conservation) relevant to the treaty;
 - iii. Evaluate the effectiveness and performance of management strategies; and
 - iv. Evaluate the effectiveness of alternative regulatory and production strategies recommended by the Parties.
2. When the Parties provide stock composition information for fisheries, the Committee shall evaluate and report its conclusions using bilaterally agreed upon methods.
3. Canada and the United States shall assess catch levels and make attempts to collect additional genetic samples from any Chum salmon caught during the July 1 through September 15 time period in the Boundary Area fisheries (U.S. Areas 4B, 5, 6C, 7 and 7A; Canadian Areas 18, 19, 20, 21, and 29).
4. During the period from July 1 through September 15, Canada will require the live release of Chum salmon from all purse seine gear fishing in the Strait of Juan de Fuca (Canadian Area 20) and the United States will require the same for the non-Indian seine fisheries in Areas 7 and 7A. Note: By U.S. regulation, purse seine fisheries are not permitted in U.S. Areas 4B, 5 and 6C.

5. Canada will manage its Johnstone Strait, Strait of Georgia, and Fraser River Chum salmon fisheries to provide continued rebuilding of depressed naturally spawning Chum salmon stocks, and, to the extent practicable, not increase interceptions of U.S. origin Chum salmon. Terminal fisheries conducted on specific stocks with identified surpluses will be managed to minimize interception of non-targeted stocks.
6. Canada will manage its Johnstone Strait mixed stock fishery as follows:
 - a) Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical;
 - b) For run sizes above the critical threshold, Canada will conduct fisheries with an exploitation rate of up to 20% in Johnstone Strait of Inside Southern Chum salmon; and
 - c) When run sizes are expected to be below the critical threshold, Canada will notify the United States and will only conduct assessment fisheries and non-commercial fisheries. Commercial fisheries targeting Chum salmon will be suspended.
7. Canada will manage its Fraser River fisheries for Chum salmon as follows:
 - a) For Fraser River terminal area run sizes, identified in-season, at abundance levels lower than 900,000 Chum salmon, the Canadian commercial Chum salmon fisheries within the Fraser River and in associated marine areas (Area 29), will be suspended; and
 - b) For Fraser River terminal area run sizes, identified in-season at levels greater than 900,000 Chum salmon, Canadian commercial Chum salmon fisheries within the Fraser River, shall be guided by the limits of the in-river Total Allowable Catch set by Canada.
8. Canada will manage the Nitinat gill net and purse seine fisheries for Chum salmon to minimize the harvest of non-targeted stocks.
9. Canada shall conduct a genetic sampling program of Chum salmon taken in the West Coast Vancouver Island troll fishery if early-season catch information indicates that catch totals for the July 1 through September 15 season may reach levels similar to 1985 and 1986. Sampling, should it occur, will include catches taken from the southern areas (Canadian Areas 121-124).
10. The United States will manage its Chum salmon fishery in Areas 7 and 7A as follows:
 - a) Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical;
 - b) For run sizes below the critical threshold, the U.S. catch of Chum salmon in Areas 7 and 7A shall be limited to Chum salmon taken incidentally to other species and in other minor fisheries, but shall not exceed 20,000, provided that catches for the purpose of genetic stock identification sampling shall not be included in the aforementioned limit;

- c) For run sizes above the critical threshold, the base catch ceiling for the U.S. Chum salmon fisheries in Areas 7 and 7A will be 130,000 Chum salmon;
 - d) Canada will provide a run size estimate of Chum salmon entering the Fraser River no later than October 22. If the estimate is less than 900,000, the U.S. will limit its fishery impacts on Fraser River Chum salmon by restricting catch in Areas 7 and 7A to not exceed 20,000 additional Chum from the day following the date the U.S. is notified. The total catch is not to exceed the catch ceiling of 130,000 Chum salmon;
 - e) U.S. commercial fisheries for fall Chum salmon in Areas 7 and 7A will not occur prior to October 10;
 - f) The U.S. will manage the Areas 7 and 7A fisheries for Chum salmon with the intent to minimize the harvest of non-targeted species;
 - g) No U.S. catch shortfalls may be accrued; however any overages shall be carried forward as indicated in (h) and (i);
 - h) Due to management imprecision, a catch in the U.S. of up to 135,000 Chum salmon will not result in an overage calculation. Catches in excess of 135,000 Chum salmon shall result in an overage being calculated by subtracting 130,000 from the total Chum catch. Overages will be accounted for by reducing the U.S. annual catch ceiling in up to two subsequent non-critical Inside Southern Chum salmon years; and
 - i) From the day following the date the U.S. is notified of a run size below the critical threshold as defined in 10(b) or (d), any catches in excess of 20,000 Chum salmon will result in an overage. Overages will be accounted for by reducing the U.S. annual catch ceilings in up to two subsequent non-critical Inside Southern Chum salmon years.
11. The United States shall conduct its Chum salmon fishery in the Strait of Juan de Fuca (United States Areas 4B, 5, and 6C) so as to maintain the limited effort nature of this fishery, and, to the extent practicable, not increase interceptions of Canadian origin Chum salmon. The United States shall continue to monitor this fishery to determine if recent catch levels indicate an increasing level of interception.
 12. All information concerning by-catch of other salmon species from the Chum salmon fisheries covered by this chapter will be shared between the Parties in the annual Post Season Report.
 13. Should circumstances arise that are inconsistent with either Parties understanding of the intent of the chapter, the Southern Panel will discuss the matter postseason and explore options for taking the appropriate corrective actions.

Appendix B-1. United States commercial salmon catch areas for the Strait of Juan de Fuca and Northern Puget Sound.



Appendix B-2. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Marine areas



Appendix B.1: Canadian Pacific Railway Management Areas (MAs) by Division (D) - Lower Fraser River and surrounding areas



2002 NBCE Test Content Blueprint

(Based on the 2002 NBCE Examination)

Overview

General Information

- (1) Test Name/Version
- (2) Test Description
 - (a) Report on the conditions of the test
- (3) Test Content
 - (a) Major content of the examination report
 - (b) The content of the test report is consistent with the content of the 2002 NBCE Test Content Blueprint
- (4) Test content is of general knowledge, ability, and skill
 - (a) Knowledge, ability, and skill is of the test
 - (b) Test content is of the test
 - (c) Test content is of the test
- (5) Test of the test is of the test
- (6) Test of the test is of the test
- (7) Test of the test is of the test
- (8) Test of the test is of the test
- (9) Test of the test is of the test
- (10) Test of the test is of the test
- (11) Test of the test is of the test
- (12) Test of the test is of the test
- (13) Test of the test is of the test
- (14) Test of the test is of the test
- (15) Test of the test is of the test
- (16) Test of the test is of the test
- (17) Test of the test is of the test
- (18) Test of the test is of the test
- (19) Test of the test is of the test
- (20) Test of the test is of the test
- (21) Test of the test is of the test
- (22) Test of the test is of the test
- (23) Test of the test is of the test
- (24) Test of the test is of the test
- (25) Test of the test is of the test
- (26) Test of the test is of the test
- (27) Test of the test is of the test
- (28) Test of the test is of the test
- (29) Test of the test is of the test
- (30) Test of the test is of the test
- (31) Test of the test is of the test
- (32) Test of the test is of the test
- (33) Test of the test is of the test
- (34) Test of the test is of the test
- (35) Test of the test is of the test
- (36) Test of the test is of the test
- (37) Test of the test is of the test
- (38) Test of the test is of the test
- (39) Test of the test is of the test
- (40) Test of the test is of the test
- (41) Test of the test is of the test
- (42) Test of the test is of the test
- (43) Test of the test is of the test
- (44) Test of the test is of the test
- (45) Test of the test is of the test
- (46) Test of the test is of the test
- (47) Test of the test is of the test
- (48) Test of the test is of the test
- (49) Test of the test is of the test
- (50) Test of the test is of the test
- (51) Test of the test is of the test
- (52) Test of the test is of the test
- (53) Test of the test is of the test
- (54) Test of the test is of the test
- (55) Test of the test is of the test
- (56) Test of the test is of the test
- (57) Test of the test is of the test
- (58) Test of the test is of the test
- (59) Test of the test is of the test
- (60) Test of the test is of the test
- (61) Test of the test is of the test
- (62) Test of the test is of the test
- (63) Test of the test is of the test
- (64) Test of the test is of the test
- (65) Test of the test is of the test
- (66) Test of the test is of the test
- (67) Test of the test is of the test
- (68) Test of the test is of the test
- (69) Test of the test is of the test
- (70) Test of the test is of the test
- (71) Test of the test is of the test
- (72) Test of the test is of the test
- (73) Test of the test is of the test
- (74) Test of the test is of the test
- (75) Test of the test is of the test
- (76) Test of the test is of the test
- (77) Test of the test is of the test
- (78) Test of the test is of the test
- (79) Test of the test is of the test
- (80) Test of the test is of the test
- (81) Test of the test is of the test
- (82) Test of the test is of the test
- (83) Test of the test is of the test
- (84) Test of the test is of the test
- (85) Test of the test is of the test
- (86) Test of the test is of the test
- (87) Test of the test is of the test
- (88) Test of the test is of the test
- (89) Test of the test is of the test
- (90) Test of the test is of the test
- (91) Test of the test is of the test
- (92) Test of the test is of the test
- (93) Test of the test is of the test
- (94) Test of the test is of the test
- (95) Test of the test is of the test
- (96) Test of the test is of the test
- (97) Test of the test is of the test
- (98) Test of the test is of the test
- (99) Test of the test is of the test
- (100) Test of the test is of the test

Test content is of the test

- (1) Test content is of the test

Question 1: Multiple Choice (10 points)

Question 1: Multiple Choice (10 points)

- a) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- b) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- c) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- d) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?

Question 2: Multiple Choice (10 points)

- a) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- b) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- c) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?
- d) The following is a list of the top 10 most popular movies in the United States in 2010. Which of the following is the correct order of the movies from highest to lowest popularity?

Appendix D Concept proposals (see work plans) submitted to Pacific Salmon Commission Southern Endowment Fund

Project Title: Joint U.S. and C.A. Mixed-stock Chum Fisheries Sampling Design

In order to facilitate management responses to Southern Chum stock strength, in accordance with Annex IV, Chapter 6 of the Pacific Salmon Treaty (Treaty) it is necessary to provide the catch composition in fisheries targeting Southern origin Chum populations. We are proposing to sample Southern BC and U.S. mixed stock Chum fisheries to determine stock composition to the Canadian Conservation Unit (CU) and United States Management Unit (MU) level using genetic mixed stock analysis. In addition, run timing, distribution and diversion by CU and MU will be explored for run reconstruction. This proposal is informed by the results from the Southern fund project (Southern Study Area Chum stock distribution assessment in Washington and San Juan Island, Point Roberts and BC southern Gulf fisheries) that occurred 2007–2010. The goal is to sample the mixed stock Chum fisheries for four consecutive years in a multi-agency sampling effort.

Project Title: Chum Salmon Southern Area Genetic Baseline Enhancement (year 2)

This project enables the short- and long-term goals of the Joint Chum Technical Committee (Chum TC) to support the development, evaluation, and improvement of microsatellite and SNP data for identification of southern stocks in mixed-stock fisheries (page 3 of the Chum TC Joint Directions white paper, January, 2010). We propose an updated survey of genetic variation in representative population samples of Chum salmon from southern British Columbia (SBC) and Washington State (WA) to enhance the Chum salmon population genetic baseline for management and research. Tissue samples will be collected from representative stocks (roughly 30 stocks for initial evaluation) and genotyped at microsatellite (mSAT) and single nucleotide polymorphism (SNP) loci.


This study will enhance the existing mSAT baseline, particularly in the U.S., and thus will provide more accurate estimates in its present day use in international fishery estimates. This study will also enhance the existing SNP baseline to improve accuracy in international estimates. The enhanced SNP baseline will build on the panel currently being developed for Chum salmon originating from British Columbia and Washington in an ongoing project of UW and WDFW. The mSAT and SNP baselines will be compared for resolving power individually and in combination (concatenate data types) for mixed stock fishery and other fishery management questions through simulations and analyses of known and unknown mixed samples.

The primary application of these genetic data is resolving the stock composition of mixed stock fisheries. Genetic baselines will also be used to develop a new fisheries model that incorporates genetic and environmental information to manage fisheries. Other analyses could include relative reproductive success of hatchery and wild-origin spawners and the impact of supplementation programs, identifying origins of juvenile Chum salmon using coastal nursery areas and address other management questions. The data set will also be evaluated for its utility in identification of southern stocks harvested in other areas, such as the Bering Sea trawl fishery.

Project Title: Development of Chum Genetic and Environmental Management (ChumGEM) Model: Phase I

The Chum Technical Committee (ChumTC), in consideration of the requirements of the latest version of Annex IV, Chapter 6 (Chum Annex) of the Pacific Salmon Treaty, has determined that a significant amount of stock assessment work should be undertaken by the parties, in order to provide the level of information necessary for the successful implementation of the Annex. ChumTC has developed a strategic plan which has been accepted by the Pacific Salmon Commission (PSC). From this plan the ChumTC submitted three proposals in 2012 which were approved by the Southern Fund Committee. The approved proposals were for building and expanding genetic baselines using genetic stock identification (GSI) techniques, coordinating and conducting a GSI workshop, and coordinating and conducting mixed stock fisheries sample collections and analysis. The Southern Panel of the PSC has identified the first phase of the Chum Genetic and Environmental Management model (ChumGEM) as a top research priority and is looking for proposals to accomplish the first phase of this model development.

A key outcome of the Chum Strategic Plan is a model that incorporates output from Tier 1 components (GSI baseline development and GSI analysis of fishery collections) with the Tier 2 information including fishery and escapement data as well as environmental components. Phase I entails compiling, standardizing, and formatting the appropriate input data, and developing a beta version of the model. The first phase of the model will be used for post-season run reconstructions for Southern BC and Washington Chum salmon. Subsequent phases will incorporate modules for forecasting and annual fishery planning, and long term stock management. The first phase of this model development will be focused on a module to address key post-season requirements under the Chum Annex.



The development of this module will incorporate the following inputs:

1. Escapement by Management Unit (MU) and Conservation Unit (CU)
2. Catch of Chum salmon spatially and temporally
3. Catch composition by MU and CU based on Genetic Stock Identification
4. Biological data (i.e. age, sex and length) from both escapement and fishery samples